

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, 15/62, 15/81, 5/12, 1/19, C07K 14/715, 16/18, C12Q 1/68		A1	(11) International Publication Number: WO 95/33051
(21) International Application Number: PCT/US95/06639		(43) International Publication Date: 7 December 1995 (07.12.95)	
(22) International Filing Date: 25 May 1995 (25.05.95)		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(30) Priority Data: 08/250,858 27 May 1994 (27.05.94) US 08/331,394 28 October 1994 (28.10.94) US 08/446,915 22 May 1995 (22.05.95) US		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(71) Applicant: GENENTECH, INC. [US/US]; 460 Point San Bruno Boulevard, South San Francisco, CA 94080-4990 (US).			
(72) Inventors: GOEDDEL, David, V.; 2115 Forestview, Hillsborough, CA 94011 (US). ROTHE, Mike; 801 North Humboldt Street #406, San Mateo, CA 94401 (US).			
(74) Agents: DREGER, Ginger, R. et al.; Genentech, Inc., 460 Point San Bruno Boulevard, South San Francisco, CA 94080-4990 (US).			

(54) Title: TUMOR NECROSIS FACTOR RECEPTOR-ASSOCIATED FACTORS

(57) Abstract

The invention concerns new tumor necrosis factor receptor associated factors, designated TRAFs. The new factors are capable of specific association with the intracellular domain of the type 2 TNF receptor (TNF-R2) and CD40, and are involved in the mediation of TNF and CD40 ligand biological activities.

TRAF2 1 MAAAGVTPGSELELLOPQFSELTLLQVLEAKVLCSEAKHILRRPFOAGCG

TRAF2 21 HRYCEPCLTSLISGGPQNCAGGVSELYSEGISILESSAAPPDAAAREV

TRAF2 261 SLPATVCPNDQCTWSTLKEVESCHNEGLPPFLTECPACKOLVRLSEKH

TRAF1 1MASSRAPDENEFQFQCPFA

TRAF2 121 HTEQCEPKELLSCHNCRAPCENYDLEVHYEVCPKFPITCDQCGKKKIPRE

TRAF1 20 PCDDPSEPNVLOCTACLSENLDDEDRIOPECRADNMPVSPN SPLTGE

TRAF2 221 TFODNTRACSKORVLDMFTVGCSEMYSTENLDHQLQALREHLALLLS

TRAF1 20 EVMSDN...AAEINMPFAGVGCSEKQSPQSMDEHATSOSSNLYLLAV

TRAF2 222 FLISAGAPQTLNOVSEHLQRI.....

TRAF1 122 LEWESEPPQNLGGAPMALSENLSLQLOAAVSEATQDLEVDCYRAPCEE

TRAF2 272QQLERQIATPSNIVDFLRSEVAVYTAELCSRNH

TRAF1 126 QEELALONLVEKLLAQSEHLRYDAMIVAVLREVEASHLALALASINDS

TRAF2 226 RLDQKIEALSNVQQLERSIGLQLANADLQKRVSELYSTYDGVPIHWK

TRAF1 226 QLDRELLSEQRNVSELDOTLQKQDVLOKLSNLSLMLSEASFQDTANE

TRAF2 228 IDDFIRKCHAYAGRTPAIFSPAFYTSRYGYEMLCLVYLNQDQITGRGTHL

TRAF1 244 ITNVIRKCHSEVCDRTVSLSPAFYTSRYGYEMLCLVYLNQDQITGRGTHL

TRAF2 428 SLFFVYVKEPNDALLQMPFHQKVTLLLDHMRREHVIDAFAPDVTSSPQ

TRAF1 228 SLFIIVIRDEVDALLQMPFANQVTLLLDHMRREHVIDAFAPDVTSSPQ

TRAF2 428 RPIVSDNIIASGCPFLCPVSKME.AKDSVYDDAIFIXAIVDTGL

TRAF1 244 RPIQDETIVASGCPFLFLSKQSPKATVYHDTNLSKQIVDTSA

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TUMOR NECROSIS FACTOR RECEPTOR-ASSOCIATED FACTORS

Field of the Invention

The present invention concerns novel polypeptide factors. More particularly, the invention concerns factors associated with the type 2 tumor necrosis factor receptor (TNF-R2).

Background of the Invention

5 Tumor necrosis factor (TNF, also referred to as TNF- α) is a potent cytokine produced mainly by activated macrophages and a few other cell types. The large number of biological effects elicited by TNF include hemorrhagic necrosis of transplanted tumors, cytotoxicity, a role in endotoxin shock, inflammatory, immunoregulatory, proliferative, and antiviral responses [reviewed in Goeddel, D.V. *et al.*, Cold Spring Harbor Symposia on Quantitative Biology 51, 597-609 (1986); Beutler, B. and Cerami, A., Ann. Rev. Biochem. 57, 505-518 (1988); Old, L.J., Sci. Am. 258(5), 59-75 (1988); Fiers, W. FEBS Lett. 285(2), 199-212 (1991)]. The literature has reported that TNF and other cytokines such as IL-1 may protect against the deleterious effects of ionizing radiation produced during the course of radiotherapy, such as denaturation of enzymes, lipid peroxidation, and DNA damage [(Neta *et al.*, J. Immunol. 136(7): 2483, (1987); Neta *et al.*, Fed. Proc. 46: 1200 (abstract), (1987); Urbaschek *et al.*, Lymphokine Res. 6: 179 (1987); U.S. Patent No. 4,861,587; Neta *et al.*, J. Immunol. 140: 108 (1988)]. A related molecule, lymphotoxin (LT, also referred to as TNF- β), that is produced by activated lymphocytes shows a similar but not identical spectrum of biological activities as TNF (see, e.g. Goeddel, D.V. *et al.*, *supra*, and Fiers, W., *supra*). TNF was described by Pennica *et al.*, Nature 312, 721 (1984); LT was described by Gray *et al.*, Nature 312, 724 (1984).

20 The first step in the induction of the various cellular responses mediated by TNF or LT is their binding to specific cell surface receptors. Two distinct TNF receptors of approximately 55-kDa (TNF-R1) and 75-kDa (TNF-R2) have been identified [Hohmann, H.P. *et al.*, J. Biol. Chem. 264, 14927-14934 (1989); Brockhaus, M. *et al.*, Proc. Natl. Acad. Sci. USA 87, 3127-3131 (1990)]; and human and mouse cDNAs corresponding to both receptor types have been isolated and characterized [Loetscher, H. *et al.*, Cell 61, 351 (1990); Schall, T.J. *et al.*, Cell 61, 361 (1990); Smith, C.A. *et al.*, Science 248, 1019 (1990); Lewis, M. *et al.*, Proc. Natl. Acad. Sci. USA 88, 2830-2834 (1991); Goodwin, R.G. *et al.*, Mol. Cell. Biol. 11, 3020-3026 (1991)]. Both TNF-Rs share the typical structure of cell surface receptors including extracellular, transmembrane and intracellular regions. The extracellular portions of both receptors are found naturally also as soluble TNF-binding proteins [Nophar, Y. *et al.*, EMBO J. 9, 3269 (1990); and Kohno, T. *et al.*, Proc. Natl. Acad. Sci. U.S.A. 87, 8331 (1990)]. The amino acid sequence of human TNF-R1 and the underlying nucleotide sequence are disclosed in EP 417,563 (published 20 March 1991), whereas EP 418,014 (published 20 March 1991) discloses the amino acid and nucleotide sequences of human TNF-R2.

Although not yet systematically investigated, the majority of cell types and tissues appear to express both TNF receptors.

35 The individual roles of the two TNF receptors, and particularly those of TNF-R2, in cell signaling are far from entirely understood, although studies performed by poly- and monoclonal antibodies (mAbs) that

specific for either TNF-R1 or TNF-R2 have provided some very valuable insight into the functions and interactions of these receptors.

It has been observed that both polyclonal and monoclonal antibodies directed against TNF-R1 can act as specific agonists for this receptor and elicit several TNF activities such as cytotoxicity, fibroblast proliferation, resistance to chlamydiae, and synthesis of prostaglandin E₂ [Engelmann, H. *et al.*, J. Biol. Chem. 265, 14497-14504 (1990); Espevik, T. *et al.*, J. Exp. Med. 171, 415-426 (1990); Shalaby, M.R. *et al.*, J. Exp. Med. 172, 1517-1520 (1990)].

In addition, polyclonal antibodies to both murine TNF-R1 and TNF-R2 have been developed, have been shown to behave as specific receptor agonists and induce a subset of murine TNF activities. While the murine TNF-R1 was shown to be responsible for signaling cytotoxicity and the induction of several genes, the murine TNF-R2 was shown to be capable of signaling proliferation of primary thymocytes and a cytotoxic T cell line, CT6 [Tartaglia, L.A. *et al.*, Proc. Natl. Acad. Sci. USA 88, 9292-9296 (1991)]. The ability of TNF-R2 to stimulate human thymocyte proliferation has been demonstrated in experiments with monoclonal antibodies directed against the human receptor.

Monoclonal antibodies against human TNF-R1 that block the binding of TNF to TNF-R1 and antagonize several of the TNF effects have also been described [Espevik, T. *et al.*, *supra*; Shalaby, M.R. *et al.*, *supra*; Naume, B. *et al.*, J. Immunol. 146, 3035-3048 (1991)].

In addition, several reports described monoclonal antibodies directed against TNF-R2 that can partially antagonize the same TNF responses (such as cytotoxicity and activation of NF- κ B) that are induced by TNF-R1 agonists [Shalaby, M.R. *et al.*, *supra*; Naume, B. *et al.*, *supra*; and Hohmann, H.P. *et al.*, J. Biol. Chem. 265, 22409-22417 (1990)].

It is now well established that although the two human TNF receptors are both active in signal transduction, they are able to mediate distinct cellular responses. While TNF-R1 appears to be responsible for signaling most TNF responses, the thymocyte proliferation stimulating activity of TNF is specifically mediated by TNF-R2. In addition, both TNF-R1 and TNF-R2 have been shown to independently mediate the activation of the transcription factor NF- κ B by TNF [Lenardo & Baltimore, Cell 58, 227-229 (1989); Læg Reid, A. *et al.*, J. Biol. Chem. 269, 7785-7791 (1994); Rothe *et al.*, Cell 78, 681-692 (1994); Wiegmann *et al.*, J. Biol. Chem. 267, 17997-18001 (1992)]. NF- κ B is a member of the Rel family of transcriptional activators that control the expression of a variety of important cellular and viral genes [Lenardo & Baltimore, *supra*, and Thanos and Maniatis, Cell 80, 529-532 (1995)]. TNF-R2 also mediates the transcriptional induction of the granulocyte-macrophage colony stimulating factor (GM-CSF) gene [Miyatake *et al.*, EMBO J. 4, 2561-2568 (1985); Stanley *et al.*, EMBO J. 4, 2569-2573 (1985)] and the A20 zinc finger protein gene [Opipari *et al.*, J. Biol. Chem. 265, 14705-14708 (1990)] in CT6 cells, and participates as an accessory component to TNF-R1 in the signaling of responses primarily mediated by TNF-R1, like cytotoxicity [Tartaglia, L.A. and Goeddel, D.V., Immunol. Today 13, 151-153 (1992)].

Summary of the Invention

Although TNF itself, the TNF receptors and TNF activities mediated by the two receptors have been studied extensively, the post-receptor signal transduction mechanisms are unknown [see the review article by Beyaert, R. & Fiers, W., "Molecular mechanisms of tumor necrosis factor-induced cytotoxicity: what we do understand and what we do not", FEBS Letters 340, 9-16 (1994)]. This is especially true for the very first step in the TNF receptor signal transduction cascade, i.e. for the question of how the membrane-bound receptor sends a signal into the cell after activation by the ligand, TNF.

The present invention is based on the hypothesis that polypeptide factors associated with the intracellular domain of TNF-R2 exist and participate in the TNF-R2 signal transduction cascade. More specifically, this invention is based on research directed to the identification and isolation of native polypeptide factors that are capable of association with the intracellular domain of TNF-R2 and participate in the intracellular post-receptor signaling of TNF biological activities.

It is known that the TNF induced proliferation of murine CT6 cells is mediated by TNF-R2 [Tartaglia *et al.*, (1991), *supra*]. To identify factors that are associated with the intracellular domain of hTNF-R2, the receptor was immunoprecipitated from lysates of [³⁵S]-labeled transfected CT6 cells and from unlabeled transfected human embryonic kidney 293 cells, which were then incubated with labeled lysate from untransfected CT6 cells. Several polypeptides with apparent molecular weights of about 45 to 50-56 kD and one with an approximate molecular weight of 68 kD were specifically coprecipitated with the immunoprecipitated hTNF-R2. These are hereinafter collectively referred to as tumor necrosis factor receptor associated polypeptides, or TRAFs. Of the factors identified two have so far been purified and cloned. These two factors are designated as tumor necrosis factor receptor associated factors 1 and 2 (TRAF1 and TRAF2; SEQ. ID. NOs: 2 and 4). A comparison of the amino acid sequences of TRAF1 and TRAF2 revealed that they share a high degree of amino acid identity in their C-terminal domains (53% identity over 230 amino acids), while their N-terminal domains are unrelated. These new factors are believed to play a key role in the post-receptor signaling of TNF. Since the intracellular domain of TNF-R2 does not display any sequence homology to any other known receptor or protein, these signaling molecules might represent a novel signal transduction mechanism, the understanding of which can greatly contribute to the development of new strategies to improve the therapeutic value of TNF.

In one aspect, the present invention concerns a family of novel factors (TRAFs) capable of specific association with the intracellular domain of a native TNF-R2. The invention specifically concerns tumor necrosis factor receptor associated factors 1 and 2 (TRAF1 and TRAF2, SEQ. ID. NOs. 2 and 4), including the native factors from any human or non-human animal species and their functional derivatives.

In another aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence encoding a TRAF polypeptide.

In yet another aspect, the invention concerns an expression vector comprising the foregoing nucleic acid molecule operably linked to control sequences recognized by a host cell transformed with the vector.

In a further aspect, the invention concerns a host cell transformed with the foregoing expression vector.

In a still further aspect, the invention concerns molecules (including polypeptides, e.g. antibodies and TRAF analogs and fragments, peptides and small organic molecules) which disrupt the interaction of a TNF-R2 receptor associated factor and TNF-R2.

The invention specifically concerns antibodies, capable of specific binding to a native TRAF polypeptide, and hybridoma cell lines producing such antibodies.

In a different aspect, the invention concerns a method of using a nucleic acid molecule encoding a TRAF polypeptide as hereinabove defined, comprising expressing such nucleic acid molecule in a cultured host cell transformed with a vector comprising said nucleic acid molecule operably linked to control sequences recognized by the host cell transformed with the vector, and recovering the encoded polypeptide from the host cell.

The invention further concerns a method for producing a TRAF polypeptide as hereinabove defined, comprising inserting into the DNA of a cell containing nucleic acid encoding said polypeptide a transcription modulatory element in sufficient proximity and orientation to the nucleic acid molecule to influence the transcription thereof.

The invention also provides a method of determining the presence of a TRAF polypeptide, comprising hybridizing DNA encoding such polypeptide to a test sample nucleic acid and determining the presence of TRAF polypeptide DNA.

In a further aspect, the invention concerns an isolated nucleic acid molecule encoding a fusion of an intracellular domain sequence of a native TNF-R2 and the DNA-binding domain of a transcriptional activator.

In a still further aspect, the invention concerns an isolated nucleic acid molecule encoding a fusion of a TRAF to the activation domain of a transcriptional activator.

The invention further concerns hybrid (fusion) polypeptides encoded by the foregoing nucleic acids.

The invention also covers vectors comprising one or both of the nucleic acid molecules encoding the foregoing fusion proteins.

In a different aspect, the invention concerns an assay for identifying a factor capable of specific binding to the intracellular domain of a native TNF-R2, comprising

(a) expressing, in a single host cell carrying a reporter gene, nucleic acid molecules encoding a polypeptide comprising a fusion of an intracellular domain sequence of a native TNF-R2 to the DNA-binding domain of a transcriptional activator, and a fusion of a candidate factor to the activation domain of a transcriptional activator; and

(b) monitoring the binding of the candidate factor to the TNF-R2 intracellular domain sequence by detecting the molecule encoded by the reporter gene.

The invention further relates to an assay for identifying a factor capable of specific association with the intracellular domain of a native TNF-R2, comprising

(a) expressing nucleic acid molecules encoding a polypeptide comprising a fusion of an intracellular domain sequence of a native TNF-R2 to the DNA-binding domain of a transcriptional activator, and a second polypeptide comprising a fusion of a candidate polypeptide factor to the activation domain of

a transcriptional activator, in a single host cell transfected with nucleic acid encoding a polypeptide factor capable of specific binding to said TNF-R2, and with nucleic acid encoding a reporter gene; and

(b) monitoring the association of said candidate factor with said TNF-R2 or with said polypeptide factor capable of specific binding to said TNF-R2 by detecting the polypeptide encoded by said reporter gene.

In a further aspect, the invention concerns an assay for identifying a molecule capable of disrupting the association of a TRAF with the intracellular domain of a native TNF-R2, comprising contacting a cell expressing 1. a fusion of an intracellular domain sequence of a native TNF-R2 to the DNA-binding domain of a transcriptional activator, 2. a fusion of a native TRAF polypeptide to the activation domain of said transcriptional activator, and 3. a reporter gene, with a candidate molecule, and monitoring the ability of said candidate molecule to disrupt the association of said TRAF and TNF-R2 intracellular domain sequence by detecting the molecule encoded by the reporter gene. The cell, just in the previous assays is preferably a yeast cell.

In addition to the "two-hybrid" format described above, the assay may be performed in any conventional binding/inhibitor assay format. For example, one binding partner (TNF-R2 or TRAF) may be immobilized, and contacted with the other binding partner equipped with a detectable label, such as a radioactive label, e.g. ^{32}P and the binding (association) of the two partners is detected in the presence of a candidate inhibitor. The design of a specific binding assay is well within the skill of a person skilled in the art.

In a different aspect, the invention concerns a method of amplifying a nucleic acid test sample comprising priming a nucleic acid polymerase reaction with nucleic acid encoding a TRAF polypeptide, as defined above.

In another aspect, the invention concerns a method for detecting a nucleic acid sequence coding for a polypeptide molecule which comprises all or part of a TRAF polypeptide or a related nucleic acid sequence, comprising contacting the nucleic acid sequence with a detectable marker which binds specifically to at least part of the nucleic acid sequence, and detecting the marker so bound.

In yet another aspect, the invention concerns a method for treating a pathological condition associated with a TNF biological activity mediated, fully or partially, by TNF-R2, comprising administering to a patient in need a therapeutically effective amount of a TRAF or a molecule capable of disrupting the interaction of a TRAF and TNF-R2.

In a still further embodiment, the invention concerns an assay for identifying an inhibitor of the interaction of a TRAF protein with CD40 comprising contacting recombinant host cells coexpressing a TRAF protein capable of direct or indirect binding CD40, CD40 and a reporter gene the expression of which is dependent on the CD40:TRAF interaction, with candidate inhibitors and selecting a molecule which inhibits the expression of said reporter gene.

In yet another embodiment, the invention concerns an assay for identifying an inhibitor of the interaction of a TRAF protein with LMP1 comprising contacting recombinant host cells coexpressing a TRAF protein capable of direct or indirect binding of LMP1, LMP1 and a reporter gene the expression of which is

dependent on the LMP1:TRAF interaction, with candidate inhibitors and selecting a molecule which inhibits the expression of said reporter gene.

Just like the assay based upon the interaction of TRAF proteins with TNF-R2, the CD40:TRAF and LMP1:TRAF assays are preferable performed in the "two-hybrid" format. However, they may alternatively
5 be performed in any conventional binding/inhibitor assay format, just like those mentioned before. The design of a specific assay format is well within the skill of an ordinary artisan.

The invention further concerns TRAF2(87-501) or a functional derivative thereof capable of inhibiting a biological activity mediated by TNF-R2, CD40 or LMP1.

Brief Description of Drawings

10 **Figure 1.** Activation of the transcription factor NF- κ B through TNF-R2 in CT6 cells.
6 μ g of nuclear extract prepared from CT6 cells that had been stimulated for 20 min with a 1:500 dilution of anti-mTNF-R2 polyclonal antibodies or the respective preimmune serum were incubated with a radiolabeled double-stranded oligonucleotide containing either two wild-type (wt) or mutant (mt) NF- κ B binding sites and analyzed for the induction of NF- κ B DNA-binding activity by electrophoretic mobility shift assay [Schütze
15 *et al.*, Cell **71**, 765-776 (1992)]. Binding reactions were either performed without competitor oligonucleotides or in the presence of a 500 fold excess of unlabeled competitor oligonucleotides containing mutant NF- κ B binding sites or a binding site for the transcription factor AP-1 [Angel, P. *et al.*, Mol. Cell. Biol. **7**: 2256-2266 (1987)]. F and B refer to free oligonucleotide probe and oligonucleotide probe in a complex with protein, respectively.

20 **Figure 2.** Immunoprecipitation of hTNF-R2.
(A) 35 S-labeled CT6 cells or CT6 cells expressing the hTNF-R2 were stimulated for 10 min with 100 ng/ml hTNF or left untreated. The cells were lysed and the hTNF-R2 immunoprecipitated as described in the text and analyzed by SDS-PAGE and autoradiography. The asterisk marks the band corresponding to the 75-80 kd hTNF-R2.

25 (B) The hTNF-R2 was immunoprecipitated from unstimulated or TNF-stimulated 293 or 293/TNF-R2 cells and incubated with lysates from 35 S-labeled CT6 cells. Arrows indicate bands of 45 to 50-56 kd and 68 kd that coprecipitate specifically with the hTNF-R2 in both experiments. Molecular weight markers are indicated on the right in kd.

Figure 3. Purification of GST-hTNF-R2_{icd} fusion protein.
30 Glutathione-S-transferase (GST) and GST-hTNF-R2_{icd} fusion protein were expressed in *E. coli*, purified as described in the text and analyzed by SDS-PAGE and Coomassie staining. Molecular weight markers are indicated on the right in kd.

Figure 4. Coprecipitation of GST-hTNF-R2_{icd} fusion protein in CT6 cell extracts.
GST and GST-hTNF-R2_{icd} fusion protein beads were incubated with lysates from 35 S-labeled CT6 cells as
35 described in the text and analyzed by SDS-PAGE and autoradiography. Arrows indicate bands of 45 to 50-56 kd and 68 kd that coprecipitate specifically with the GST-hTNF-R2_{icd} fusion protein. Molecular weight markers are indicated on the right in kd.

Figure 5. Coprecipitation of GST-mutant hTNF-R2_{icd} fusion proteins in CT6 cell extracts.

GST and GST-fusion proteins containing mutant intracellular domains of the hTNF-R2 were coupled to glutathione-agarose beads, incubated with lysates from ³⁵S-labeled CT6 cells and analyzed by SDS-PAGE and autoradiography. Arrows indicate bands of 45 to 50-56 kd and 68 kd that coprecipitate specifically with the GST-fusion proteins containing the wild type (wt), the mutant -16, the Δ304-345 and the 384-424 intracellular domains of hTNF-R2 but are not associated with the mutant -37 and -59 intracellular domains. Note that the pattern of these bands is compressed in some cases due to the unlabeled fusion proteins migrating at the same size. Molecular weight markers are indicated on the right in kd.

Figure 6. Competition of TNF-R2 associated factors with GST-hTNF-R2_{icd} fusion proteins.

(A) The hTNF-R2 was immunoprecipitated from 293 and 293/TNF-R2 cells and incubated with lysates from ³⁵S-labeled CT6 cells that had been preincubated with 50 μl of the indicated GST-hTNF-R2_{icd} fusion protein beads as competitor. Reactions were analyzed by SDS-PAGE and autoradiography. Arrows indicate bands of 45 to 50-56 kd and 68 kd that coprecipitate specifically with the hTNF-R2 and that are depleted by preincubation with GST-fusion proteins containing the wild type (wt) and the mutant -16 intracellular domains of hTNF-R2 but not by preincubation with the mutant -37 and -59 intracellular domain fusion proteins.

(B) The 68 kd region of a similar experiment as described in (A) is shown.

Molecular weight markers are indicated on the right in kd.

Figure 7. Coprecipitation of GST-hTNF-R2_{icd} fusion protein in Jurkat cell extracts.

GST and GST-hTNF-R2_{icd} fusion protein beads were incubated with lysates from ³⁵S-labeled Jurkat cells that had been stimulated for 10 min with 100 ng/ml hTNF or left untreated. Reactions were analyzed by SDS-PAGE and autoradiography. Arrows indicate bands of 45 to 50-56 kd, 67 kd and 73-75 kd that coprecipitate specifically with the GST-hTNF-R2_{icd} fusion protein. Molecular weight markers are indicated on the right in kd.

Figure 8. Subcellular localization of TNF-R2 associated factors.

Cytoplasmic and cell membrane fractions were prepared from ³⁵S-labeled CT6 cells as described in the text. These fractions and a detergent (total) extract from CT6 cells were incubated with GST and GST-hTNF-R2_{icd} fusion beads, and the reactions analyzed by SDS-PAGE and autoradiography. Arrows indicate bands of 45 to 50-56 kd and 68 kd that coprecipitate specifically with the GST-hTNF-R2_{icd} fusion protein. Molecular weight markers are indicated on the right in kd.

Figure 9. Purification of TNF-R2 associated factors.

Large scale purification of TNF-R2 associated factors from CT6 cells by GST-hTNF-R2_{icd} fusion protein affinity chromatography was performed as described in the text. One tenth of the obtained material was analyzed by SDS-PAGE and silver staining. Arrows indicate bands of 45 to 50-56 kd and 68-70 kd that were eluted specifically from the GST-hTNF-R2_{icd} fusion protein affinity column. Molecular weight markers are indicated on the right in kd.

Figure 10. Nucleotide and predicted amino acid sequence of the TRAF1 cDNA (SEQ. ID. NOS: 1 and 2).

The nucleic acid sequence of the TRAF1 cDNA is shown with numbering starting from the first base after the *Sall* cloning linker. The deduced protein sequence is displayed above with numbering from the initiation methionine. In-frame stop codons upstream of the initiation methionine are underlined. Amino acids identified by sequencing the purified TRAF1 protein are indicated in bold. The TRAF domain (see text) comprises amino acids 180 (>) -409 (<). The potential leucine zipper region (see text) extends between amino acids 183 (+) -259 (-). Amino acids within this region defining the heptade motif are indicated in italic.

Figure 11. Nucleotide and predicted amino acid sequence of the TRAF2 cDNA (SEQ. ID. NOS: 3 and 4).

The nucleic acid sequence of the longest TRAF2 cDNA is shown with numbering starting from the first base after the *Sall* cloning linker. In addition, the first nucleotide of four independently isolated pPC86 cDNA inserts (*) and the longest λ phage cDNA insert (^) is indicated. The deduced protein sequence is displayed above with numbering from the putative initiation methionine, which is in-frame with the GAL4 activation domain coding region in all isolated pPC86TRAF2 cDNA clones (see text). Cysteine and histidine residues defining the RING finger motif and the two TFIIIA-like zinc finger motifs (see text) are indicated in bold or underlined, respectively. The TRAF domain (see text) comprises amino acids 272 (>) -501 (<). The potential leucine zipper region (see text) extends between amino acids 275 (+) -351 (-). Amino acids within this region defining the heptade motif are indicated in italic.

Figure 12. Sequence similarity of regions in TRAF2 to zinc-binding motifs.

(A) Comparison of amino acid sequences containing RING finger motifs. The TRAF2 RING finger motif is aligned with the respective zinc-binding motifs of the regulatory protein COPI from *A. thaliana* [Deng *et al.*, Cell **71**, 791-801 (1992); SEQ. ID. NO: 5], the human estrogen-responsive finger protein EFP [Inoue *et al.*, Proc. Natl. Acad. Sci. USA **90**, 11117-11121 (1993); SEQ. ID. NO: 6], the RAD18 and UVS-2 gene products required for DNA repair in *S. cerevisiae* and *N. crassa*, respectively [Jones *et al.*, Nucl. Acids Res. **16**, 7119-7131 (1988); SEQ. ID. NO: 7]; [Tomita *et al.*, Mol. Gen. Genet. **238**, 225-233 (1993); SEQ. ID. NO: 8], the human V(D)J recombination activating gene product RAG-1 [Schatz *et al.*, Cell **59**, 1035-1048 (1989); SEQ. ID. NO: 9], the human 52 kd ribonucleoprotein SS-A/Ro [Chan *et al.*, J. Clin. Invest. **87**, 68-76 (1987)]; Itoh *et al.*, J. Clin. Invest. **87**, 177-186 (1987); A⁵² in ref. 1 is P⁵² in ref. 2; SEQ. ID. NO: 10], human RING1 [Lovering, GBTRANS accession number Z14000 (1992); SEQ. ID. NO: 11], mouse T lymphocyte regulatory protein RPT-1 [Patarca *et al.*, Proc. Natl. Acad. Sci. USA **85**, 2733-2737 (1988); SEQ. ID. NO: 12], human regulatory protein RFP [Takahashi *et al.*, Mol. Cell. Biol. **8**, 1853-1856 (1988); SEQ. ID. NO: 13], and the product of the human proto-oncogen c-cbl [Blake *et al.*, Oncogene **6**, 653-657 (1991); SEQ. ID. NO: 14].

(B) Comparison of amino acid sequences containing TFIIIA-type zinc finger motifs. A region in TRAF2 comprising two contiguous repeats of the consensus sequence C_H-X₂₋₄-C_H-X₂₋₁₅-C_H-X₂₋₄-C_H [Berg, J. Biol. Chem. **265**, 6513-6516 (1990)] is aligned with similar zinc-binding motifs of the developmentally regulated DG17 gene product from *D. discoideum* [Driscoll & Williams, Mol. Cell. Biol. **7**, 4482-4489 (1987); SEQ. ID. NO: 15], the transcription factor IIIA form *X. laevis* [Miller *et al.*, EMBO J. **4**, 1609-1614 (1985); SEQ. ID. NO: 16], the *Xenopus* zinc finger proteins XLCOF14 and XFIN [Nietfeld *et al.*, J.

Mol. Biol. **208**, 639-659 (1989); SEQ. ID. NO: 17]; [Ruiz i Altaba *et al.*, EMBO J. **6**, 3065-3070 (1987); SEQ. ID. NO: 18], the mouse ZFY1/2 and MFG2 gene products [Mardon & Page, Cell **56**, 765-770 (1989); SEQ. ID. NO: 19]; [Passananti *et al.*, Proc. Natl. Acad. Sci. USA **86**, 9421-9471 (1989); SEQ. ID. NO: 20], and the RAD18 and UVS-2 proteins (see above; SEQ. ID. NOS: 21 and 22).

5 **Figure 13. Homology between TRAF1 and TRAF2.**

An optimized alignment of the protein sequences of TRAF1 and TRAF2 is shown. Identical amino acids are boxed. The C-terminal TRAF domain (see text) comprises amino acids 180-409 of TRAF1 and 272-501 of TRAF2.

Figure 14. Hydropathy analysis of TRAF1 and TRAF2.

10 Hydropathy profiles of the amino acid sequences of TRAF1 (A) and TRAF2 (B) were obtained by the method of Kyte and Doolittle, J. Mol. Biol. **157**, 105-132 (1982) using a window of twenty amino acids. The numbers under each plot indicate positions of the amino acids of the respective protein.

Figure 15. Northern blot analysis of TRAF1 and TRAF2 mRNA.

(A) Northern blot analysis of TRAF1 and TRAF2 mRNA in CT6 cells. There is 3 µg of
15 poly(A)⁺RNA from CT6 cells per lane.

(B) Northern blot analysis of TRAF1 and TRAF2 mRNA in mouse tissues. Mouse multiple tissue northern blots (Clontech) were hybridized with radiolabeled TRAF1 and TRAF2 probes as described in the text.

Figure 16. Coprecipitation of GST-TRAF2 fusion protein in 293 cell extracts.

20 GST and GST-TRAF2 fusion protein beads were incubated with lysates from 293 and 293/TNF-R2 cells as described in the text. Reactions were analyzed by SDS-PAGE and Western blot analysis using anti-human TNF-R1 monoclonal antibody 986 (0.5 µg/ml) and anti-human TNF-R2 monoclonal antibody 1036 (0.5 µg/ml). An arrow indicates the 75-80 kd hTNF-R2 band that is coprecipitated specifically with the GST-TRAF2 fusion protein. Molecular weight markers are indicated on the right in kd.

25 **Figure 17. Transcription factor NF-κB is activated by TRAF2 overexpression.** 293 cells (10⁶) were transfected with 7.5 µg of pRK control (lanes 1-5) or expression vectors for TRAF1 (lane 6), TRAF3 (lane 7), and TRAF2 (lanes 7-16). The expression vectors and transient transfection in 293 cells are described in Example 6. Cells were treated with 100 ng/ml human TNF (lane 2), pre-immune serum (lane 3), anti-human TNF-R1 antibodies (lane 4) or anti-human TN F-R2 antibodies (lane 5) at a dilution of 1:500 for 1 h prior
30 to harvest [Rothe *et al.*, Cell **78**, 681 (1994)]. Nuclear extracts were prepared 24 hours after transfection, and 6 µg aliquots were incubated with a radiolabeled double-stranded oligonucleotide containing two NF-κB-binding sites [Schütze *et al.*, Cell **71**, 765 (1992)]. Individual reactions were supplemented with a 50-fold excess of unlabeled competitor oligonucleotide containing either wild-type (lane 9) or mutated (lane 10) NF-κB sequence [Schütze *et al.* supra]. Reaction mixtures were incubated with 1 µl of pre-immune serum (lane
35 11), anti-p50 serum (lane 12), anti-p65 serum (lane 13), anti-c-rel serum (lane 14), anti-relB serum (lane 15) or anti-p52 serum (lane 16) [Santa Cruz Biotechnology]. F and B refer to free oligonucleotide probe or oligonucleotide probe in a complex with protein, respectively.

Figure 18. Overexpression of TRAF2 induces NF- κ B dependent reporter gene activity. (A) Effect of TRAF overexpression on NF- κ B dependent reporter gene activity in 293 cells. 293 cells were transiently cotransfected with an E-selectin-luciferase reporter gene plasmid [Schindler and Baichwal, *Mol. Cell. Biol.* 14: 5820 (1994)] and TRAF expression vectors (1 μ g) as indicated in Example 6. Cells were either untreated (closed bars) or stimulated for 8 h with 100 ng/ml human TNF (hatched bars) prior to harvest. Luciferase activities were determined and normalized based on β -galactosidase expression. Values shown are averages (mean \pm SD) of one representative experiment in which each transfection was performed in triplicate. (B) Effect of TRAF expression on NF- κ B dependent reporter gene activity in CT6 cells. CT6 cells were transiently cotransfected with an E-selectin-luciferase reporter gene plasmid (as described in Example 6) and TRAF expression vectors (6 μ g). (Transient transfection of CT6 cells was performed using the DEAE-dextran method [F.M. Ausubell *et al.*, Current Protocols in Molecular Biology (Green Publishing Associates/Wiley & Sons, Inc.) New York, (1994)]. 10⁷ CT6 cells were transfected with a total of 10 μ g of plasmid DNA in 250 μ g/ml DEAE-dextran for 90 min. Reporter gene activity was assayed 40 h after transfection). After 24 h cells were left untreated (closed bars) or stimulated for an additional 16 h with 100 ng/ml murine TNF (hatched bars) prior to harvest. Luciferase activities were determined and normalized based on β -galactosidase expression. Values shown are averages (mean \pm SD) of one representative experiment in which each transfection was performed in triplicate. Numbers represent the fold induction of reporter gene activity by TNF stimulation for the respective transfections.

Figure 19. TRAF2 mediates induction of NF- κ B dependent reporter gene activity by TNF-R2 and CD40. (A) Effect of coexpression of TNF-R2 and TRAFs on NF- κ B dependent reporter gene activity in 293 cells. 293 cells were transiently cotransfected with an E-selectin-luciferase reporter gene plasmid [Schindler and Baichwal, *supra*] and expression vectors for TNF-R2 (1 μ g) and TRAFs (1 μ g) as described in Example 6. Cells were either untreated (closed bars) or stimulated for 8 h with 100 ng/ml human TNF (hatched bars) prior to harvest. Luciferase activities were determined and normalized based on β -galactosidase expression. Values shown are averages (mean \pm SD) of one representative experiment in which each transfection was performed in triplicate. (B) Effect of coexpression of CD40 and TRAFs on NF- κ B dependent reporter gene activity in 293 cells. 293 cells were transiently cotransfected with an E-selectin-luciferase reporter gene plasmid [Schindler and Baichwal, *supra*] and expression vectors for CD40 (1 μ g) and TRAFs (1 μ g) as indicated in Example 6. After 24 h, cells were harvested and luciferase activities determined and normalized (closed bars). Values shown are averages (mean \pm SD) of one representative experiment in which each transfection was performed in triplicate.

Detailed Description of the Invention

A. Definitions

The phrases "factor," "tumor necrosis factor receptor associated factor"; "TNF-R2 associated factor" and "TRAF" are used interchangeably and refer to a native factor capable of specific association with the intracellular domain of a native TNF-R2, and functional derivatives of such native factor. In the context of this definition the phrase "specific association" is used in the broadest sense, and includes direct binding to

a site or region within the intracellular domain of a native TNF-R2 of the human or of any animal species, and indirect association with a native TNF-R2 intracellular domain, mediated by a further molecule, such as another TRAF. The phrase "native TRAF" designates a TRAF polypeptide as occurring in nature in any cell type of any human or non-human animal species, with or without the initiating methionine, whether purified
5 from native source, synthesized, produced by recombinant DNA technology or by any combination of these and/or other methods. Native TRAFs specifically include monomeric, homo- and heterodimeric and homo- and heterooligomeric forms of such naturally occurring polypeptides. The native murine TRAF1 and TRAF2 polypeptides (SEQ. ID. NOS: 2 and 4) are unrelated in the region of the N-terminal region (RING finger domain) of TRAF2. In contrast, they exhibit a 41% sequence identity outside of this domain. This homology
10 is subdivided into two regions. A region of low sequence identity of 28% comprises amino acids 28-136 of TRAF1 and amino acids 159-271 of TRAF2. This region is separated by a 43 amino acid insertion in TRAF1 from a region of high sequence identity of 53% over 230 amino acids comprising the C-terminal domains of TRAF1 (amino acids 180-409) and TRAF2 (amino acids 272-501). This sequence similarity provides evidence of a novel structural domain that is hereby designated the TRAF domain. The native TRAF
15 polypeptides preferably share a novel sequence motif in the C-terminal portion of their amino acid sequences, and preferably are at least about 40%, more preferably at least about 50%, most preferably at least about 55% homologous within this C-terminal "TRAF domain". The "TRAF domain" encompasses about amino acids 272 to 501 of the native mouse TRAF2 amino acid sequence, about amino acids 180 to 409 of the native mouse TRAF1 amino acid sequence, and homologous domains of other native TRAFs and their functional
20 derivatives. The native TRAFs further have the preferred property of binding CD40 and/or LMP1 and mediating their biological activities.

The terms "native type 2 TNF receptor" and "native TNF-R2" are used interchangeably, and refer to any naturally occurring (native) type 2 TNF receptor from any (human and non-human) animal species, with or without the initiating methionine and with or without a signal sequence attached to the N-terminus,
25 whether purified from native source, synthesized, produced by recombinant DNA technology or by any combination of these and/or other methods.

The terms "native human type 2 TNF receptor" and "native human TNF-R2", which are used interchangeably, refer to a human TNF-R2 having the amino acid sequence disclosed in EP 418,014 (published 20 March 1991), with or without the initiating methionine and with or without a signal sequence
30 attached to the N-terminus, whether purified from native source, synthesized, produced by recombinant DNA technology or by any combination of these and/or other methods, and other naturally occurring human TNF-R2 variants, including soluble and variously glycosylated forms of native full-length human TNF-R2, whether purified from natural sources, synthetically produced in vitro or obtained by genetic manipulation including methods of recombinant DNA technology.

35 A "functional derivative" of a native polypeptide is a compound having a qualitative biological activity in common with the native polypeptide. Thus, a functional derivative of a native TRAF polypeptide is a compound that has a qualitative biological activity in common with a native TRAF. "Functional derivatives" include, but are not limited to, fragments of native polypeptides from any animal species

(including humans), and derivatives of native (human and non-human) polypeptides and their fragments, provided that they have a biological activity in common with a respective native polypeptide. "Fragments" comprise regions within the sequence of a mature native polypeptide. The term "derivative" is used to define amino acid sequence and glycosylation variants, and covalent modifications of a native polypeptide, whereas the term "variant" refers to amino acid sequence and glycosylation variants within this definition. Preferably, the functional derivatives are polypeptides which have at least about 65% amino acid sequence identity, more preferably about 75% amino acid sequence identity, even more preferably at least about 85% amino acid sequence identity, most preferably at least about 95% amino acid sequence identity with the sequence of a corresponding native polypeptide. Most preferably, the functional derivatives of a native TRAF polypeptide retain or mimic the region or regions within the native polypeptide sequence that directly participate in the association with the TNF-R2 intracellular domain and/or in homo- or heterodimerization. The phrase "functional derivative" specifically includes peptides and small organic molecules having a qualitative biological activity in common with a native TRAF.

The term "biological activity" in the context of the definition of functional derivatives is defined as the possession of at least one adhesive, regulatory or effector function qualitatively in common with a native polypeptide (e.g. TRAF). A preferred biological property of the functional derivatives of the native TRAF polypeptides herein is their ability to associate with the intracellular domain of a native TNF-R2 (either by direct binding or via interaction with another TRAF), and thereby mediate or block a biological response signaled (exclusively or partially) by the TNF-R2 with which they are associated. Another preferred biological activity is the ability of the TRAF polypeptides of the present invention to signal the activation of the transcription factor NF- κ B. Further preferred functional derivatives of native TRAF proteins are characterized by blocking TNF-R2 mediated NF- κ B activation, while typically retaining the ability to interact with the cytoplasmic domain of TNF-R2 as well as with other TRAFs. Another preferred biological activity is the ability of certain TRAF functional derivatives to associate with the cytoplasmic domain of CD40. Such functional derivatives can mediate or block the biological activities of CD40. A further preferred group of TRAF functional derivatives is capable of associating with the LMP1 oncogene (see Example 6) and mediate or block its biological activity.

"Identity" or "homology" with respect to a native polypeptide and its functional derivative is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues of a corresponding native polypeptide, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent homology, and not considering any conservative substitutions as part of the sequence identity. Neither N- or C-terminal extensions nor insertions shall be construed as reducing identity or homology. Methods and computer programs for the alignment are well known in the art.

The TRAF polypeptides of the present invention specifically include native murine TRAF1 (SEQ. ID. NO: 2) and native murine TRAF2 (SEQ. ID. NO: 4) their homo- and heterodimeric and homo- and heterooligomeric forms, and their analogs in other mammalian species, such as rat, porcine, equine, cow, higher primates, and humans, and the functional derivatives of such native polypeptides. The functional derivatives of a native TRAF1 or native TRAF2 receptor are preferably encoded by DNAs capable, under

stringent conditions, of hybridizing to the complement of a DNA encoding a native TRAF polypeptide. More preferably, the functional derivatives share at least about 40% sequence homology, more preferably at least about 50% sequence homology, even more preferably at least about 55% sequence homology, most preferably at least about 60% sequence homology with any domain, and preferably with the TNF-R2 binding domain(s) and/or the dimerization domain(s), of a native TRAF polypeptide. In a most preferred embodiment, a functional derivative will share at least about 50% sequence homology, more preferably at least about 55% sequence homology, most preferably at least about 60% sequence homology with the C-terminal TRAF region of murine TRAF2, or are encoded by DNA capable of hybridizing, under stringent conditions, with the complement of DNA encoding the TRAF region of murine TRAF2. In another preferred embodiment, a functional derivative of a native TRAF polypeptide is a fragment having a "TRAF" domain as the only functionally intact domain, wherein the "TRAF" domain encompasses about amino acids 272 to 501 of the native mouse TRAF2 amino acid sequence, about amino acids 180 to 409 of the native mouse TRAF1 amino acid sequence, and homologous domains of other native TRAFs. In a further preferred embodiment, a functional derivative of a native TRAF polypeptide lacks the RING finger domain of TRAF2. One of the latter variants, TRAF2(87-501), is completely defective in signalling NF- κ B activation.

The "stringent conditions" are overnight incubation at 42 °C in a solution comprising: 20% formamide, 5xSSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 μ g/ml denatured, sheared salmon sperm DNA.

The terms "amino acid" and "amino acids" refer to all naturally occurring L- α -amino acids. The amino acids are identified by either the single-letter or three-letter designations:

Asp	D	aspartic acid	Ile	I	isoleucine
Thr	T	threonine	Leu	L	leucine
Ser	S	serine	Tyr	Y	tyrosine
Glu	E	glutamic acid	Phe	F	phenylalanine
Pro	P	proline	His	H	histidine
Gly	G	glycine	Lys	K	lysine
Ala	A	alanine	Arg	R	arginine
Cys	C	cysteine	Trp	W	tryptophan
Val	V	valine	Gln	Q	glutamine
Met	M	methionine	Asn	N	asparagine

These amino acids may be classified according to the chemical composition and properties of their side chains. They are broadly classified into two groups, charged and uncharged. Each of these groups is divided into subgroups to classify the amino acids more accurately:

I. Charged Amino Acids

Acidic Residues: aspartic acid, glutamic acid

Basic Residues: lysine, arginine, histidine

II. Uncharged Amino Acids

5 Hydrophilic Residues: serine, threonine, asparagine, glutamine

Aliphatic Residues: glycine, alanine, valine, leucine, isoleucine

Non-polar Residues: cysteine, methionine, proline

Aromatic Residues: phenylalanine, tyrosine, tryptophan

10 The term "amino acid sequence variant" refers to molecules with some differences in their amino acid sequences as compared to a native amino acid sequence.

Substitutional variants are those that have at least one amino acid residue in a native sequence removed and a different amino acid inserted in its place at the same position. The substitutions may be single, where only one amino acid in the molecule has been substituted, or they may be multiple, where two or more amino acids have been substituted in the same molecule.

15 Insertional variants are those with one or more amino acids inserted immediately adjacent to an amino acid at a particular position in a native sequence. Immediately adjacent to an amino acid means connected to either the α -carboxy or α -amino functional group of the amino acid.

20 Deletional variants are those with one or more amino acids in the native amino acid sequence removed. Ordinarily, deletional variants will have one or two amino acids deleted in a particular region of the molecule.

25 The term "glycosylation variant" is used to refer to a glycoprotein having a glycosylation profile different from that of a native counterpart or to glycosylated variants of a polypeptide unglycosylated in its native form(s). Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side-chain of an asparagine residue. The tripeptide sequences, asparagine-X-serine and asparagine-X-threonine, wherein X is any amino acid except proline, are recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be involved in O-linked glycosylation.

30 Antibodies (Abs) and immunoglobulins (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules which lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas.

35 Native antibodies and immunoglobulins are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number

of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains [Clothia *et al.*, J. Mol. Biol. **186**, 651-663 (1985); Novotny and Haber, Proc. Natl. Acad. Sci. USA **82**, 4592-4596 (1985)].

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed through the variable domains of antibodies. It is concentrated in three segments called complementarity determining regions (CDRs) or hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen binding site of antibodies [see Kabat, E.A. *et al.*, Sequences of Proteins of Immunological Interest, National Institute of Health, Bethesda, MD (1991)]. The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

Papain digestion of antibodies produces two identical antigen binding fragments, called Fab fragments, each with a single antigen binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen recognition and binding site. This region consists of a dimer of one heavy and one light chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen binding site on the surface of the V_H - V_L dimer. Collectively, the six CDRs confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other, chemical couplings of antibody fragments are also known.

The light chains of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g. IgG-1, IgG-2, IgG-3, and IgG-4; IgA-1 and IgA-2. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called α , delta, epsilon, γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The term "antibody" is used in the broadest sense and specifically covers single monoclonal antibodies (including agonist and antagonist antibodies), antibody compositions with polyepitopic specificity, as well as antibody fragments (e.g., Fab, F(ab')₂, and Fv), so long as they exhibit the desired biological activity.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler & Milstein, Nature 256:495 (1975), or may be made by recombinant DNA methods [see, e.g. U.S. Patent No. 4,816,567 (Cabilly *et al.*)].

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity [U.S. Patent No. 4,816,567; Cabilly *et al.*; Morrison *et al.*, Proc. Natl. Acad. Sci. USA 81, 6851-6855 (1984)].

"Humanized" forms of non-human (e.g. murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂, or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-

human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibody may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details see: Jones *et al.*, Nature **321**, 522-525 [1986]; Reichmann *et al.*, Nature **332**, 323-329 [1988]; and Presta, Curr. Op. Struct. Biol. **2** 593-596 [1992]).

In the context of the present invention the expressions "cell", "cell line", and "cell culture" are used interchangeably, and all such designations include progeny. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological property, as screened for in the originally transformed cell, are included.

"Transformation" means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integration.

"Transfection" refers to the taking up of an expression vector by a host cell whether or not any coding sequences are in fact expressed.

The terms "transformed host cell" and "transformed" refer to the introduction of DNA into a cell. The cell is termed a "host cell", and it may be a prokaryotic or a eukaryotic cell. Typical prokaryotic host cells include various strains of E. coli. Typical eukaryotic host cells are mammalian, such as Chinese hamster ovary cells or human embryonic kidney 293 cells. The introduced DNA is usually in the form of a vector containing an inserted piece of DNA. The introduced DNA sequence may be from the same species as the host cell or a different species from the host cell, or it may be a hybrid DNA sequence, containing some foreign and some homologous DNA.

The terms "replicable expression vector" and "expression vector" refer to a piece of DNA, usually double-stranded, which may have inserted into it a piece of foreign DNA. Foreign DNA is defined as heterologous DNA, which is DNA not naturally found in the host cell. The vector is used to transport the foreign or heterologous DNA into a suitable host cell. Once in the host cell, the vector can replicate independently of the host chromosomal DNA, and several copies of the vector and its inserted (foreign) DNA may be generated. In addition, the vector contains the necessary elements that permit translating the foreign DNA into a polypeptide. Many molecules of the polypeptide encoded by the foreign DNA can thus be rapidly synthesized.

"Oligonucleotides" are short-length, single- or double-stranded polydeoxynucleotides that are chemically synthesized by known methods [such as phosphotriester, phosphite, or phosphoramidite chemistry, using solid phase techniques such as those described in EP 266,032, published 4 May 1988, or via

deoxynucleoside H-phosphanate intermediates as described by Froehler *et al.*, Nucl. Acids Res. **14**, 5399 (1986). They are then purified on polyacrylamide gels.

B. Identification and purification of TRAFs

The native TRAF polypeptides can be identified in and purified from certain tissues known to possess a type 2 TNF receptor (TNF-R2) mRNA and to express it at a detectable level. Thus, murine TRAF can, for example, be obtained from the murine interleukin 2 (IL-2)-dependent cytotoxic T cell line CT6 [Ranges *et al.* J. Immunol. **142**, 1203-1208 (1989)]. Murine TRAF1 can also be purified from spleen, lung and testis; whereas murine TRAF2 can be isolated and purified from an even larger variety of tissues, including heart, brain, spleen, lung, liver, skeletal muscle, kidney and testis (see Figure 15b). In general, TRAF proteins are expected to be expressed in human tissues that are known to express TNF-R2, although not all of such tissues will express all TRAFs. Alternatively, TRAF polypeptides can be isolated from cell lines transfected with DNA encoding a native TNF-R2 or a TNF-R2 derivative comprising intracellular domain sequences participating in the interaction with TRAFs. Factors that are associated with the intracellular domain of a native TNF-R2 can be identified by immunoprecipitation of the receptor or receptor derivative from cells expressing it. Immunoprecipitation in general consists of multiple ordered steps, including lysing the cell with detergent if the TNF-R2 is membrane-bound, binding of TNF-R2 to an anti-TNF-R2 antibody, precipitating the antibody complex, washing the precipitate, and dissociating TNF-R2 and any associated factor from the immune complex. The dissociated factor(s) can then be analyzed by electrophoretic methods. In a preferred embodiment, radiolabeled TNF-R2 (or a derivative) is immunoprecipitated with protein A-agarose (Oncogene Science) or with protein A-Sepharose (Pharmacia). In this case, the TNF-R2/anti-TNF-R2 antibody immune complexes are precipitated by *Staphylococcus aureus* protein A bound to the agarose or Sepharose. The immunoprecipitate is then analyzed by autoradiography or by fluorography, depending on the actual radiolabel used. The TRAF proteins (which are characterized by their ability to associate with the intracellular domain of TNF-R2) will coprecipitate with the receptor or receptor derivative, and can be further purified by methods known in the art, such as purification on an affinity column.

A large-scale purification scheme for purifying factors that associate with the intracellular domain of TNF-R2 takes advantage of plasmid expression vectors that direct the synthesis of foreign polypeptides in *E. coli* as fusions with the C terminus of glutathione S-transferase (GST), as described by Smith, D.B. and Johnson, K.S., Gene **67** 31-40 (1988). The intracellular domain of TNF-R2 is expressed as a fusion protein with GST in *E. coli* recombinant host cells, and can be purified from crude bacterial lysates by absorption on glutathione-agarose beads (Sigma). A cell lysate containing the factor(s) to be purified is then applied to a GST-TNF-R2 fusion protein affinity column. Protein(s) bound to the column is/are eluted, precipitated and isolated by SDS-PAGE under reducing conditions, and visualized by silver staining. GST gene fusion vectors (pGEX vectors) as well as kits for cloning and expression of GST fusion systems are commercially available from Pharmacia (see Pharmacia Catalog, 1994, pages 133; and 142-143).

Purified protein can be either sequenced directly by automated Edman degradation with a model 470A Applied Biosystems gas phase sequencer equipped with a 120A PTH amino acid analyzer or sequenced

after digestion with various chemicals or enzymes. PTH amino acids were integrated using a ChromPerfect data system (Justice Innovations, Palo Alto, CA). Sequence interpretation can be performed on a VAX 11/785 Digital Equipment Corporation computer as described by Henzel *et al.*, J. Chromatography 404, 41 (1987). In some cases, eluates electrophoresed on SDS polyacrylamide gels are electrotransferred to a PVDF
5 membrane (ProBlott, ABI, Foster City, CA) and stained with Coomassie Brilliant Blue R250 (Sigma). The specific protein is excised from the blot for N-terminal sequencing. To determine internal protein sequences, purified fractions obtained by reverse phase capillary HPLC are typically dried under vacuum (SpeedVac), resuspended in appropriate buffers, and digested with cyanogen bromide, and/or various proteases, such as trypsin, the lysine-specific enzyme Lys-C (Wako Chemicals, Richmond, VA) or Asp-N (Boehringer
10 Mannheim, Indianapolis, Ind.). After digestion, the resultant peptides are sequenced as a mixture or are resolved by HPLC.

C. Recombinant production of TRAF polypeptides

Preferably, the TRAF polypeptides are prepared by standard recombinant procedures by culturing cells transfected to express TRAF polypeptide nucleic acid (typically by transforming the cells with an
15 expression vector) and recovering the polypeptide from the cells. However, it is envisioned that the TRAF polypeptides may be produced by homologous recombination, or by recombinant production methods utilizing control elements introduced into cells already containing DNA encoding an TRAF polypeptide. For example, a powerful promoter/enhancer element, a suppressor, or an exogenous transcription modulatory element may be inserted in the genome of the intended host cell in proximity and orientation sufficient to influence the
20 transcription of DNA encoding the desired TRAF polypeptide. The control element does not encode the TRAF polypeptide, rather the DNA is indigenous to the host cell genome. One next screens for cells making the polypeptide of this invention, or for increased or decreased levels of expression, as desired.

Thus, the invention contemplates a method for producing a TRAF polypeptide comprising inserting into the genome of a cell containing nucleic acid encoding a TRAF polypeptide a transcription modulatory
25 element in sufficient proximity and orientation to the nucleic acid molecule to influence transcription thereof, with an optional further step of culturing the cell containing the transcription modulatory element and the nucleic acid molecule. The invention also contemplates a host cell containing the indigenous TRAF polypeptide nucleic acid molecule operably linked to exogenous control sequences recognized by the host cell.

1. Isolation of DNA encoding the TRAF polypeptides

For the purpose of the present invention, DNA encoding a TRAF polypeptide can be obtained from
30 cDNA libraries prepared from tissue believed to possess a type 2 TNF receptor (TNF-R2) mRNA and to express it at a detectable level. For example, cDNA library can be constructed by obtaining polyadenylated mRNA from a cell line known to express TNF-R2, and using the mRNA as a template to synthesize double stranded cDNA. Human and non-human cell lines suitable for this purpose have been listed hereinabove.
35 It is noted, however, that TNF-R2 is known to be expressed in a large variety of further tissues which can all potentially serve as a source of TRAF cDNA, even though not all members of the TRAF family will be

expressed in all TNF-R2 expressing tissues. Alternatively, DNA encoding new TRAF polypeptides can be obtained from cDNA libraries prepared from tissue known to express a previously identified TRAF polypeptide at a detectable level. The TRAF polypeptide genes can also be obtained from a genomic library, such as a human genomic cosmid library.

5 Libraries, either cDNA or genomic, are screened with probes designed to identify the gene of interest or the protein encoded by it. For cDNA expression libraries, suitable probes include monoclonal and polyclonal antibodies that recognize and specifically bind to a TRAF polypeptide. For cDNA libraries, suitable probes include carefully selected oligonucleotide probes (usually of about 20-80 bases in length) that encode known or suspected portions of a TRAF polypeptide from the same or different species, and/or
10 complementary or homologous cDNAs or fragments thereof that encode the same or a similar gene. Appropriate probes for screening genomic DNA libraries include, without limitation, oligonucleotides, cDNAs, or fragments thereof that encode the same or a similar gene, and/or homologous genomic DNAs or fragments thereof. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures as described in Chapters 10-12 of Sambrook *et al.*, Molecular Cloning: A Laboratory Manual,
15 New York, Cold Spring Harbor Laboratory Press, 1989).

A preferred method of practicing this invention is to use carefully selected oligonucleotide sequences to screen cDNA libraries from various tissues. The oligonucleotide sequences selected as probes should be sufficient in length and sufficiently unambiguous that false positives are minimized. The actual nucleotide sequence(s) is/are usually designed based on regions of a TRAF which have the least codon redundancy. The
20 oligonucleotides may be degenerate at one or more positions. The use of degenerate oligonucleotides is of particular importance where a library is screened from a species in which preferential codon usage is not known.

The oligonucleotide must be labeled such that it can be detected upon hybridization to DNA in the library being screened. The preferred method of labeling is to use ATP (e.g., $\gamma^{32}\text{P}$) and polynucleotide kinase
25 to radiolabel the 5' end of the oligonucleotide. However, other methods may be used to label the oligonucleotide, including, but not limited to, biotinylation or enzyme labeling.

cDNAs encoding TRAFs can also be identified and isolated by other known techniques of recombinant DNA technology, such as by direct expression cloning or by using the polymerase chain reaction (PCR) as described in U.S. Patent No. 4,683,195, issued 28 July 1987, in section 14 of Sambrook *et al.*,
30 Molecular Cloning: A Laboratory Manual, second edition, Cold Spring Harbor Laboratory Press, New York, 1989, or in Chapter 15 of Current Protocols in Molecular Biology, Ausubel *et al.* eds., Greene Publishing Associates and Wiley-Interscience 1991. This method requires the use of oligonucleotide probes that will hybridize to DNA encoding a TRAF.

According to a preferred method for practicing the invention, the coding sequences for TRAF
35 proteins can be identified in a recombinant cDNA library or a genomic DNA library based upon their ability to interact with the intracellular domain of a TNF-R2. For this purpose one can use the yeast genetic system described by Fields and co-workers [Fields and Song, Nature (London) **340**, 245-246 (1989); Chien *et al.*, Proc. Natl. Acad. Sci. USA **88**, 9578-9582 (1991)] as disclosed by Chevray and Nathans [Proc. Natl. Acad.

Sci. USA 89, 5789-5793 (1991)]. Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, while the other one functioning as the transcription activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

To directly isolate genes encoding proteins that associate with the intracellular domain of TNF-R2, DNA encoding a TNF-R2 intracellular domain or a fragment thereof is cloned into a vector containing DNA encoding the DNA-binding domain of GAL4. A plasmid cDNA library is then constructed by cloning double-stranded cDNA encoding a candidate factor in a vector comprising DNA encoding the GAL4 transcriptional activation domain. Thereafter, yeast cells containing reporter genes are cotransformed with the TNF-R2-GAL4 DNA binding domain vector and with library plasmid DNA. Typically, an *S. cerevisiae* cell containing two reporter genes: *lacZ*(β gal) and *His* genes, serves as a host for cotransformation. Yeast transformants are selected by plating on supplemented synthetic dextrose medium lacking tryptophan, leucine and histidine, and protein-protein interactions are monitored by the yeast colony filter β -galactosidase assay, essentially as described by Chevray and Nathans, *supra*. Only colonies with protein-protein interaction will grow on his⁻ plates, and are then analyzed for β -gal as a further control.

Once cDNA encoding a TRAF from one species has been isolated, cDNAs from other species can also be obtained by cross-species hybridization. According to this approach, human or other mammalian cDNA or genomic libraries are probed by labeled oligonucleotide sequences selected from known TRAF sequences (such as murine TRAF1 and TRAF2 as disclosed in the present application) in accord with known criteria, among which is that the sequence should be sufficient in length and sufficiently unambiguous that false positives are minimized. Typically, a ³²P-labeled oligonucleotide having about 30 to 50 bases is sufficient, particularly if the oligonucleotide contains one or more codons for methionine or tryptophan. Isolated nucleic acid will be DNA that is identified and separated from contaminant nucleic acid encoding other polypeptides from the source of nucleic acid.

Once the sequence is known, the gene encoding a particular TRAF polypeptide can also be obtained by chemical synthesis, following one of the methods described in Engels and Uhlmann, *Angew. Chem. Int. Ed. Engl.* 28, 716 (1989). These methods include triester, phosphite, phosphoramidite and H-phosphonate methods, PCR and other autoprimer methods, and oligonucleotide syntheses on solid supports.

2. Amino Acid Sequence Variants of a native TRAF proteins or fragments

Amino acid sequence variants of native TRAFs and TRAF fragments are prepared by methods known in the art by introducing appropriate nucleotide changes into a native or variant TRAF DNA, or by *in vitro* synthesis of the desired polypeptide. There are two principal variables in the construction of amino acid sequence variants: the location of the mutation site and the nature of the mutation. With the exception of naturally-occurring alleles, which do not require the manipulation of the DNA sequence encoding the TRAF, the amino acid sequence variants of TRAF are preferably constructed by mutating the DNA, either to arrive at an allele or an amino acid sequence variant that does not occur in nature.

One group of the mutations will be created within the domain or domains identified as being involved in the interaction with the intracellular domain of TNF-R2, CD40 or LMP1. TRAF variants mutated to enhance their association (binding or indirect association) with TNF-R2, CD40 or LMP1 and/or to retain their binding ability while reducing or eliminating their ability to signal NF-kB activation, will be useful as inhibitors of native biological activities mediated by native TNF-R2, CD40 or LMP1 proteins. In addition, such variants will be useful in the diagnosis of pathological conditions association with the overexpression of these proteins, and in the purification of TNF-R2, CD40 or LMP1. A target for such mutations is the N-terminal RING finger domain of TRAF2 and related factors, as this domain is believed to be involved in the interaction with the intracellular domain of TNF-R2. A typical representative of such TRAF2 variants is a mutant TRAF2 protein lacking the N-terminal finger domain of native TRAF2 (TRAF2(87-501)). This variant retains the ability to interact with TNF-R2, CD40, TRAF1 and TRAF2 but is completely defective in signaling NF-kB activation. Other TRAF amino acid sequence variants which act as inhibitors of the TRAF-TNF-R2/CD40/LMP1 biological activity can, for example, be identified by the biochemical screening assays which will be described hereinbelow.

Another group of mutations will be performed within region(s) involved in interactions with other TNF-R2 associated factors. Thus, amino acid alterations within the homologous C-terminal domains (protein dimerization motif) of TRAF1, TRAF2 and other factors of the TRAF family can enhance the ability of such factors to form stable dimers which are required for signaling through the TNF-R2 receptor.

Alternatively or in addition, amino acid alterations can be made at sites that differ in TRAF proteins from various species, or in highly conserved regions, depending on the goal to be achieved.

Sites at such locations will typically be modified in series, e.g. by (1) substituting first with conservative choices and then with more radical selections depending upon the results achieved, (2) deleting the target residue or residues, or (3) inserting residues of the same or different class adjacent to the located site, or combinations of options 1-3.

One helpful technique is called "alanine scanning" [Cunningham and Wells, Science 244, 1081-1085 (1989)]. Here, a residue or group of target residues is identified and substituted by alanine or polyalanine. Those domains demonstrating functional sensitivity to the alanine substitutions are then refined by introducing further or other substituents at or for the sites of alanine substitution.

After identifying the desired mutation(s), the gene encoding a TRAF variant can, for example, be obtained by chemical synthesis as hereinabove described.

More preferably, DNA encoding a TRAF amino acid sequence variant is prepared by site-directed mutagenesis of DNA that encodes an earlier prepared variant or a nonvariant version of the TRAF. Site-directed (site-specific) mutagenesis allows the production of TRAF variants through the use of specific oligonucleotide sequences that encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 20 to 25 nucleotides in length is preferred, with about 5 to 10 residues on both sides of the junction of the sequence being altered. In general, the techniques of site-specific mutagenesis are well known in the art, as exemplified by publications such as, Edelman *et al.*, DNA **2**, 183 (1983). As will be appreciated, the site-specific mutagenesis technique typically employs a phage vector that exists in both a single-stranded and double-stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage, for example, as disclosed by Messing *et al.*, Third Cleveland Symposium on Macromolecules and Recombinant DNA, A. Walton, ed., Elsevier, Amsterdam (1981). This and other phage vectors are commercially available and their use is well known to those skilled in the art. A versatile and efficient procedure for the construction of oligodeoxyribonucleotide directed site-specific mutations in DNA fragments using M13-derived vectors was published by Zoller, M.J. and Smith, M., Nucleic Acids Res. **10**, 6487-6500 (1982)). Also, plasmid vectors that contain a single-stranded phage origin of replication [Veira *et al.*, Meth. Enzymol. **153**, 3 (1987)] may be employed to obtain single-stranded DNA. Alternatively, nucleotide substitutions are introduced by synthesizing the appropriate DNA fragment *in vitro*, and amplifying it by PCR procedures known in the art.

In general, site-specific mutagenesis herewith is performed by first obtaining a single-stranded vector that includes within its sequence a DNA sequence that encodes the relevant protein. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically, for example, by the method of Crea *et al.*, Proc. Natl. Acad. Sci. USA **75**, 5765 (1978). This primer is then annealed with the single-stranded protein sequence-containing vector, and subjected to DNA-polymerizing enzymes such as, *E. coli* polymerase I Klenow fragment, to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate host cells such as JP101 cells, and clones are selected that include recombinant vectors bearing the mutated sequence arrangement. Thereafter, the mutated region may be removed and placed in an appropriate expression vector for protein production.

The PCR technique may also be used in creating amino acid sequence variants of a TRAF. When small amounts of template DNA are used as starting material in a PCR, primers that differ slightly in sequence from the corresponding region in a template DNA can be used to generate relatively large quantities of a specific DNA fragment that differs from the template sequence only at the positions where the primers differ from the template. For introduction of a mutation into a plasmid DNA, one of the primers is designed to overlap the position of the mutation and to contain the mutation; the sequence of the other primer must be identical to a stretch of sequence of the opposite strand of the plasmid, but this sequence can be located

anywhere along the plasmid DNA. It is preferred, however, that the sequence of the second primer is located within 200 nucleotides from that of the first, such that in the end the entire amplified region of DNA bounded by the primers can be easily sequenced. PCR amplification using a primer pair like the one just described results in a population of DNA fragments that differ at the position of the mutation specified by the primer, and possibly at other positions, as template copying is somewhat error-prone.

If the ratio of template to product material is extremely low, the vast majority of product DNA fragments incorporate the desired mutation(s). This product material is used to replace the corresponding region in the plasmid that served as PCR template using standard DNA technology. Mutations at separate positions can be introduced simultaneously by either using a mutant second primer or performing a second PCR with different mutant primers and ligating the two resulting PCR fragments simultaneously to the vector fragment in a three (or more) part ligation.

In a specific example of PCR mutagenesis, template plasmid DNA (1 μ g) is linearized by digestion with a restriction endonuclease that has a unique recognition site in the plasmid DNA outside of the region to be amplified. Of this material, 100 ng is added to a PCR mixture containing PCR buffer, which contains the four deoxynucleotide triphosphates and is included in the GeneAmp^R kits (obtained from Perkin-Elmer Cetus, Norwalk, CT and Emeryville, CA), and 25 pmole of each oligonucleotide primer, to a final volume of 50 μ l. The reaction mixture is overlaid with 35 μ l mineral oil. The reaction is denatured for 5 minutes at 100°C, placed briefly on ice, and then 1 μ l Thermus aquaticus (Tag) DNA polymerase (5 units/l), purchased from Perkin-Elmer Cetus, Norwalk, CT and Emeryville, CA) is added below the mineral oil layer. The reaction mixture is then inserted into a DNA Thermal Cycler (purchased from Perkin-Elmer Cetus) programmed as follows:

2 min. 55°C,
30 sec. 72°C, then 19 cycles of the following:
30 sec. 94°C,
30 sec. 55°C, and
30 sec. 72°C.

At the end of the program, the reaction vial is removed from the thermal cycler and the aqueous phase transferred to a new vial, extracted with phenol/chloroform (50:50 vol), and ethanol precipitated, and the DNA is recovered by standard procedures. This material is subsequently subjected to appropriate treatments for insertion into a vector.

Another method for preparing variants, cassette mutagenesis, is based on the technique described by Wells *et al.* [Gene 34, 315 (1985)]. The starting material is the plasmid (or vector) comprising the TRAF DNA to be mutated. The codon(s) within the TRAF to be mutated are identified. There must be a unique restriction endonuclease site on each side of the identified mutation site(s). If no such restriction sites exist, they may be generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at appropriate locations in the TRAF DNA. After the restriction sites have been introduced into the plasmid, the plasmid is cut at these sites to linearize it. A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction site but containing the desired mutation(s) is synthesized using

standard procedures. The two strands are synthesized separately and then hybridized together using standard techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 3' and 5' ends that are compatible with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated TRAF DNA sequence.

5 Additionally, the so-called phagemid display method may be useful in making amino acid sequence variants of native or variant TRAFs or their fragments. This method involves (a) constructing a replicable expression vector comprising a first gene encoding an receptor to be mutated, a second gene encoding at least a portion of a natural or wild-type phage coat protein wherein the first and second genes are heterologous, and a transcription regulatory element operably linked to the first and second genes, thereby
10 forming a gene fusion encoding a fusion protein; (b) mutating the vector at one or more selected positions within the first gene thereby forming a family of related plasmids; (c) transforming suitable host cells with the plasmids; (d) infecting the transformed host cells with a helper phage having a gene encoding the phage coat protein; (e) culturing the transformed infected host cells under conditions suitable for forming recombinant phagemid particles containing at least a portion of the plasmid and capable of transforming the
15 host, the conditions adjusted so that no more than a minor amount of phagemid particles display more than one copy of the fusion protein on the surface of the particle; (f) contacting the phagemid particles with a suitable antigen so that at least a portion of the phagemid particles bind to the antigen; and (g) separating the phagemid particles that bind from those that do not. Steps (d) through (g) can be repeated one or more times. Preferably in this method the plasmid is under tight control of the transcription regulatory element,
20 and the culturing conditions are adjusted so that the amount or number of phagemid particles displaying more than one copy of the fusion protein on the surface of the particle is less than about 1%. Also, preferably, the amount of phagemid particles displaying more than one copy of the fusion protein is less than 10% of the amount of phagemid particles displaying a single copy of the fusion protein. Most preferably, the amount is less than 20%. Typically in this method, the expression vector will further contain a secretory signal
25 sequence fused to the DNA encoding each subunit of the polypeptide and the transcription regulatory element will be a promoter system. Preferred promoter systems are selected from *lac* Z, λ_{PL} , *tac*, T7 polymerase, tryptophan, and alkaline phosphatase promoters and combinations thereof. Also, normally the method will employ a helper phage selected from M13K07, M13R408, M13-VCS, and Phi X 174. The preferred helper phage is M13K07, and the preferred coat protein is the M13 Phage gene III coat protein. The preferred host
30 is *E. coli*, and protease-deficient strains of *E. coli*.

Further details of the foregoing and similar mutagenesis techniques are found in general textbooks, such as, for example, Sambrook *et al.*, *supra*, and *Current Protocols in Molecular Biology*, Ausubel *et al.* eds., *supra*.

Naturally-occurring amino acids are divided into groups based on common side chain properties:

- 35 (1) hydrophobic: norleucine, met, ala, val, leu, ile;
 (2) neutral hydrophobic: cys, ser, thr;
 (3) acidic: asp, glu;
 (4) basic: asn, gln, his, lys, arg;

(5) residues that influence chain orientation: gly, pro; and

(6) aromatic: trp, tyr, phe.

Conservative substitutions involve exchanging a member within one group for another member within the same group, whereas non-conservative substitutions will entail exchanging a member of one of these classes for another. Variants obtained by non-conservative substitutions are expected to result in significant changes in the biological properties/function of the obtained variant, and may result in TRAF variants which block TNF biological activities, especially if they are exclusively or primarily mediated by TNF-R2. Amino acid positions that are conserved among various species and/or various receptors of the TRAF family are generally substituted in a relatively conservative manner if the goal is to retain biological function.

Amino acid sequence deletions generally range from about 1 to 30 residues, more preferably about 1 to 10 residues, and typically are contiguous. Deletions may be introduced into regions not directly involved in the interaction with the TNF-R2 intracellular domain.

Amino acid insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions (i.e. insertions within the TRAF protein amino acid sequence) may range generally from about 1 to 10 residues, more preferably 1 to 5 residues, more preferably 1 to 3 residues. Examples of terminal insertions include the TRAF polypeptides with an N-terminal methionyl residue, an artifact of its direct expression in bacterial recombinant cell culture, and fusion of a heterologous N-terminal signal sequence to the N-terminus of the TRAF molecule to facilitate the secretion of the mature TRAF from recombinant host cells. Such signal sequences will generally be obtained from, and thus homologous to, the intended host cell species. Suitable sequences include STII or Ipp for *E. coli*, alpha factor for yeast, and viral signals such as herpes gD for mammalian cells.

Other insertional variants of the native TRAF molecules include the fusion of the N- or C-terminus of the TRAF molecule to immunogenic polypeptides, e.g. bacterial polypeptides such as beta-lactamase or an enzyme encoded by the *E. coli* trp locus, or yeast protein, and C-terminal fusions with proteins having a long half-life such as immunoglobulin regions (preferably immunoglobulin constant regions), albumin, or ferritin, as described in WO 89/02922 published on 6 April 1989.

Since it is often difficult to predict in advance the characteristics of a variant TRAF, it will be appreciated that screening will be needed to select the optimum variant. For this purpose biochemical screening assays, such as those described hereinbelow, will be readily available.

A preferred amino acid sequence variant of the native TRAF2 protein comprises amino acids 87-501 of the native protein, and is designated TRAF2(87-501). This variant, which lacks the RING finger domain, retains the ability to interact with TNF-R2, CD40 and LMP1, but is completely defective in signaling NF-kB activation.

3. Insertion of DNA into a Cloning Vehicle

Once the nucleic acid encoding a native or variant TRAF is available, it is generally ligated into a replicable expression vector for further cloning (amplification of the DNA), or for expression.

Expression and cloning vectors are well known in the art and contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. The selection of the appropriate vector will depend on 1) whether it is to be used for DNA amplification or for DNA expression, 2) the size of the DNA to be inserted into the vector, and 3) the host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the host cell for which it is compatible. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

(i) Signal Sequence Component

In general, the signal sequence may be a component of the vector, or it may be a part of the TRAF molecule that is inserted into the vector. If the signal sequence is heterologous, it should be selected such that it is recognized and processed (i.e. cleaved by a signal peptidase) by the host cell.

As the TRAF molecules are intracellular proteins, they are unlikely to have a native signal sequence. Heterologous signal sequences suitable for prokaryotic host cells are prokaryotic signal sequences, such as the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the yeast invertase, alpha factor, or acid phosphatase leaders may be used. In mammalian cell expression mammalian signal sequences are suitable.

(ii) Origin of Replication Component

Both expression and cloning vectors contain a nucleic acid sequence that enabled the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomes, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast and viruses. The origin of replication from the well-known plasmid pBR322 is suitable for most gram negative bacteria, the 2 μ plasmid origin for yeast and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Origins of replication are not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter). Most expression vectors are "shuttle" vectors, i.e. they are capable of replication in at least one class of organisms but can be transfected into another organism for expression. For example, a vector is cloned in *E. coli* and then the same vector is transfected into yeast or mammalian cells for expression even though it is not capable of replicating independently of the host cell chromosome.

DNA is also cloned by insertion into the host genome. This is readily accomplished using Bacillus species as hosts, for example, by including in the vector a DNA sequence that is complementary to a sequence found in Bacillus genomic DNA. Transfection of Bacillus with this vector results in homologous recombination with the genome and insertion of the DNA encoding the desired heterologous polypeptide. However, the recovery of genomic DNA is more complex than that of an exogenously replicated vector because restriction enzyme digestion is required to excise the encoded polypeptide molecule.

(iii) Selection Gene Component

Expression and cloning vectors should contain a selection gene, also termed a selectable marker. This is a gene that encodes a protein necessary for the survival or growth of a host cell transformed with the vector. The presence of this gene ensures that any host cell which deletes the vector will not obtain an advantage in growth or reproduction over transformed hosts. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g. ampicillin, neomycin, methotrexate or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g. the gene encoding D-alanine racemase for bacilli.

One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene express a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin [Southern *et al.*, J. Molec. Appl. Genet. 1, 327 (1982)], mycophenolic acid [Mulligan *et al.*, Science 209, 1422 (1980)], or hygromycin [Sudgen *et al.*, Mol. Cell. Biol. 5, 410-413 (1985)]. The three examples given above employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid), or hygromycin, respectively.

Other examples of suitable selectable markers for mammalian cells are dihydrofolate reductase (DHFR) or thymidine kinase. Such markers enable the identification of cells which were competent to take up the desired nucleic acid. The mammalian cell transformants are placed under selection pressure which only the transformants are uniquely adapted to survive by virtue of having taken up the marker. Selection pressure is imposed by culturing the transformants under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to amplification of both the selection gene and the DNA that encodes the desired polypeptide. Amplification is the process by which genes in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Increased quantities of the desired polypeptide are synthesized from the amplified DNA.

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium which lacks hypoxanthine, glycine, and thymidine. An appropriate host cell in this case is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity, prepared and propagated as described by Urlaub and Chasin, Proc. Nat'l. Acad. Sci. USA 77, 4216 (1980). A particularly useful DHFR is a mutant DHFR that is highly resistant to MTX (EP 117,060). This selection agent can be used with any otherwise suitable host, e.g. ATCC No. CCL61 CHO-K1, notwithstanding the presence of endogenous DHFR. The DNA encoding DHFR and the desired polypeptide, respectively, then is amplified by exposure to an agent (methotrexate, or MTX) that inactivates the DHFR. One ensures that the cell requires more DHFR (and consequently amplifies all exogenous DNA) by selecting only for cells that can grow in successive rounds of ever-greater MTX concentration. Alternatively, hosts co-transformed with genes encoding the desired polypeptide, wild-type DHFR, and another selectable marker such as the neo gene can be identified using a selection agent for the selectable marker such as G418 and then selected and amplified

using methotrexate in a wild-type host that contains endogenous DHFR. [See also U.S. Patent No. 4,965,199].

A suitable selection gene for use in yeast is the trp1 gene present in the yeast plasmid YRp7 [Stinchcomb *et al.*, Nature 282:39 (1979); Kingsman *et al.*, Gene 7:141 (1979); or Tschemper *et al.*, Gene 10:157 (1980)]. The trp1 gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics 85:12 (1977)]. The presence of the trp1 lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, Leu2 deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the Leu2 gene.

10 (iv) Promoter Component

Expression vectors, unlike cloning vectors, should contain a promoter which is recognized by the host organism and is operably linked to the nucleic acid encoding the desired polypeptide. Promoters are untranslated sequences located upstream from the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of nucleic acid under their control. They typically fall into two classes, inducible and constitutive. Inducible promoters are promoters that initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, e.g. the presence or absence of a nutrient or a change in temperature. At this time a large number of promoters recognized by a variety of potential host cells are well known. These promoters are operably linked to DNA encoding the desired polypeptide by removing them from their gene of origin by restriction enzyme digestion, followed by insertion 5' to the start codon for the polypeptide to be expressed. This is not to say that the genomic promoter for a TRAF polypeptide is not usable. However, heterologous promoters generally will result in greater transcription and higher yields of expressed TRAFs as compared to the native TRAF promoters.

Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang *et al.*, Nature 275:615 (1978); and Goeddel *et al.*, Nature 281:544 (1979)], alkaline phosphatase, a tryptophan (trp) promoter system [Goeddel, Nucleic Acids Res. 8:4057 (1980) and EPO Appln. Publ. No. 36,776] and hybrid promoters such as the tac promoter [H. de Boer *et al.*, Proc. Nat'l. Acad. Sci. USA 80:21-25 (1983)]. However, other known bacterial promoters are suitable. Their nucleotide sequences have been published, thereby enabling a skilled worker operably to ligate them to DNA encoding TRAF [Siebenlist *et al.*, Cell 20:269 (1980)] using linkers or adaptors to supply any required restriction sites. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding a TRAF.

Suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman *et al.*, J. Biol. Chem. 255:2073 (1980)] or other glycolytic enzymes [Hess *et al.*, J. Adv. Enzyme Reg. 7:149 (1978); and Holland, Biochemistry 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate

isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in R. Hitzeman *et al.*, EP 73.657A. Yeast enhancers also are advantageously used with yeast promoters.

Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CXCAAT region where X may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into mammalian expression vectors.

TRAF transcription from vectors in mammalian host cells may be controlled by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus [UK 2,211,504 published 5 July 1989], adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g. the actin promoter or an immunoglobulin promoter, from heat shock promoters, and from the promoter normally associated with the TRAF sequence, provided such promoters are compatible with the host cell systems.

The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment which also contains the SV40 viral origin of replication [Fiers *et al.*, Nature 273:113 (1978), Mulligan and Berg, Science 209, 1422-1427 (1980); Pavlakis *et al.*, Proc. Natl. Acad. Sci. USA 78, 7398-7402 (1981)]. The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment [Greenaway *et al.*, Gene 18, 355-360 (1982)]. A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in US 4,419,446. A modification of this system is described in US 4,601,978. See also, Gray *et al.*, Nature 295, 503-508 (1982) on expressing cDNA encoding human immune interferon in monkey cells; Reyes *et al.*, Nature 297, 598-601 (1982) on expressing human β -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus; Canaani and Berg, Proc. Natl. Acad. Sci. USA 79, 5166-5170 (1982) on expression of the human interferon β 1 gene in cultured mouse and rabbit cells; and Gorman *et al.*, Proc. Natl. Acad. Sci. USA 79, 6777-6781 (1982) on expression of bacterial CAT sequences in CV-1 monkey kidney cells, chicken embryo fibroblasts, Chinese hamster ovary cells, HeLa cells, and mouse HIN-3T3 cells using the Rous sarcoma virus long terminal repeat as a promoter.

(v) Enhancer Element Component

Transcription of a DNA encoding the TRAFs of the present invention by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Enhancers are relatively orientation and position independent having been found 5' [Laimins *et al.*, Proc. Natl. Acad. Sci. USA **78**, 993 (1981)] and 3' [Lasky *et al.*, Mol. Cell. Biol. **3**, 1108 (1983)] to the transcription unit, within an intron [Banerji *et al.*, Cell **33**, 729 (1983)] as well as within the coding sequence itself [Osborne *et al.*, Mol. Cell. Biol. **4**, 1293 (1984)]. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, Nature **297**, 17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the TRAF DNA, but is preferably located at a site 5' from the promoter.

15 (vi) Transcription Termination Component

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3' untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding the TRAF. The 3' untranslated regions also include transcription termination sites.

Construction of suitable vectors containing one or more of the above listed components, the desired coding and control sequences, employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required.

25 For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to transform E. coli K12 strain 294 (ATCC 31,446) and successful transformants selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion, and/or sequenced by the method of Messing *et al.*, Nucleic Acids Res. **9**, 309 (1981) or by the method of Maxam *et al.*, Methods in Enzymology **65**, 499 (1980).

30 Particularly useful in the practice of this invention are expression vectors that provide for the transient expression in mammalian cells of DNA encoding a TRAF. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the expression vector. Transient systems, comprising a suitable expression vector and a host cell, allow for the convenient positive identification of polypeptides encoded by clones DNAs, as well as for the rapid screening of such polypeptides for desired biological or physiological properties. Thus, transient expression systems are particularly useful in the invention for purposes of identifying analogs and variants of a TRAF.

Other methods, vectors, and host cells suitable for adaptation to the synthesis of the TRAF polypeptides in recombinant vertebrate cell culture are described in Getting *et al.*, Nature 293, 620-625 (1981); Mantel *et al.*, Nature 281, 40-46 (1979); Levinson *et al.*; EP 117,060 and EP 117,058. A particularly useful plasmid for mammalian cell culture expression of the TRAF polypeptides is pRK5 [EP 307,247].

5 (vii) Construction and analysis of vectors

Construction of suitable vectors containing one or more of the above listed components employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required.

For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to
10 transform *E. coli* K12 strain 294 (ATCC 31,446) and successful transformants selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion, and/or sequences by the methods of Messing *et al.*, Nuclei Acids Res. 9, 309 (1981) or by the method of Maxam *et al.*, Methods in Enzymology 65, 499 (1980).

 (viii) Transient expression vectors

15 Particularly useful in the practice of this invention are expression vectors that provide for the transient expression in mammalian cells of DNA encoding a TRAF polypeptide. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high level of a desired polypeptide encoded by the expression vector. Sambrook *et al.*, supra, pp. 16.17-16.22. Transient expression
20 systems, comprising a suitable expression vector and a host cell, allow for the convenient positive screening of such polypeptides for desired biological or physiological properties. Thus transient expression systems are particularly useful in the invention for purposes of identifying analogs and variants of native TRAF polypeptides with TRAF biological activity.

 (ix) Suitable exemplary vertebrate cell vectors

25 Other methods, vectors, and host cells suitable for adaptation to the synthesis of a TRAF polypeptide (including functional derivatives of native proteins) in recombinant vertebrate cell culture are described in Gething *et al.*, Nature 293, 620-625 (1981); Mantel *et al.*, Nature 281, 40-46 (1979); Levinson *et al.*, EP 117,060; and EP 117,058. A particularly useful plasmid for mammalian cell culture expression of a TRAF polypeptide is pRK5 (EP 307,247) or pSV16B [PCT Publication No. WO 91/08291].

30 D. Selection and Transformation of Host Cells

Suitable host cells for cloning or expressing the vectors herein are the prokaryote, yeast or higher eukaryote cells described above. Suitable prokaryotes include gram negative or gram positive organisms, for example E. coli or bacilli. A preferred cloning host is E. coli 294 (ATCC 31,446) although other gram

negative or gram positive prokaryotes such as E. coli B, E. coli X1776 (ATCC 31.537), E. coli W3110 (ATCC 27,325). Pseudomonas species. or Serratia Marcesans are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable hosts for vectors herein. Saccharomyces cerevisiae, or common baker's yeast, is the most commonly used among
 5 lower eukaryotic host microorganisms. However, a number of other genera, species and strains are commonly available and useful herein, such as S. pombe [Beach and Nurse, Nature 290, 140 (1981)], Kluyveromyces lactis [Louvencourt *et al.*, J. Bacteriol. 737 (1983)]; Yarrowia [EP 402.226]; Pichia pastoris [EP 183.070], Trichoderma reesia [EP 244.234], Neurospora crassa [Case *et al.*, Proc. Natl. Acad. Sci. USA 76, 5259-5263 (1979)]; and Aspergillus hosts such as A. nidulans [Ballance *et al.*, Biochem. Biophys. Res. Commun. 112,
 10 284-289 (1983); Tilburn *et al.*, Gene 26, 205-221 (1983); Yelton *et al.*, Proc. Natl. Acad. Sci. USA 81, 1470-1474 (1984)] and A. niger [Kelly and Hynes, EMBO J. 4, 475-479 (1985)].

Suitable host cells may also derive from multicellular organisms. Such host cells are capable of complex processing and glycosylation activities. In principle, any higher eukaryotic cell culture is workable, whether from vertebrate or invertebrate culture, although cells from mammals such as humans are preferred.
 15 Examples of invertebrate cells include plants and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as Spodoptera frugiperda (caterpillar), Aedes aegypti (mosquito), Aedes albopictus (mosquito), Drosophila melanogaster (fruitfly), and Bombux mori host cells have been identified. See, e.g. Luckow *et al.*, Bio/Technology 6, 47-55 (1988); Miller *et al.*, in Genetic Engineering, Setlow, J.K. *et al.*, eds., Vol. 8 (Plenum Publishing, 1986), pp. 277-279; and Maeda *et al.*,
 20 Nature 315, 592-594 (1985). A variety of such viral strains are publicly available, e.g. the L-1 variant of Autographa californica NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of Spodoptera frugiperda cells.

Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can be utilized as hosts. Typically, plant cells are transfected by incubation with certain strains of the bacterium Agrobacterium tumefaciens, which has been previously manipulated to contain the TRAF DNA. During incubation of the
 25 plant cell culture with A. tumefaciens, the DNA encoding a TRAF is transferred to the plant cell host such that it is transfected, and will, under appropriate conditions, express the TRAF DNA. In addition, regulatory and signal sequences compatible with plant cells are available, such as the nopaline synthase promoter and polyadenylation signal sequences. Depicker *et al.*, J. Mol. Appl. Gen. 1, 561 (1982). In addition, DNA
 30 segments isolated from the upstream region of the T-DNA 780 gene are capable of activating or increasing transcription levels of plant-expressible genes in recombinant DNA-containing plant tissue. See EP 321.196 published 21 June 1989.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) is per se well known. See Tissue Culture, Academic Press, Kruse and Patterson, editors
 35 (1973). Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney cell line [293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen. Virol. 36, 59 (1977)]; baby hamster kidney cells 9BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR [CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA 77,

4216 (1980)]; mouse sertolli cells [TM4, Mather, Biol. Reprod. 23, 243-251 (1980)]; monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells [Mather *et al.*, Annals N.Y. Acad. Sci. 383, 44068 (1982)]; MRC 5 cells; FS4 cells; and a human hepatoma cell line (Hep G2). Preferred host cells are human embryonic kidney 293 and Chinese hamster ovary cells.

Particularly preferred host cells for the purpose of the present invention are vertebrate cells producing the TRAF polypeptides.

10 In another particularly preferred embodiment, the TRAF polypeptides of the present invention are expressed in insect cells. The expression can, for example, be performed in Hi5 insect cells (Invitrogen, San Diego) using a baculovirus expression kit (BaculoGold), which is commercially available from Pharmingen, San Diego.

Host cells are transfected and preferably transformed with the above-described expression or cloning 15 vectors and cultured in conventional nutrient media modified as is appropriate for inducing promoters or selecting transformants containing amplified genes.

E. Culturing the Host Cells

Prokaryotes cells used to produced the TRAF polypeptides of this invention are cultured in suitable media as describe generally in Sambrook *et al.*, supra.

20 Mammalian cells can be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium (MEM, Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium (DMEM, Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham and Wallace, Meth. Enzymol. 58, 44 (1979); Barnes and Sato, Anal. Biochem. 102, 255 (1980), US 4,767,704; 4,657,866; 4,927,762; or 4,560,655; WO 90/03430; WO 87/00195 or US Pat. Re. 25 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and thymidine), antibiotics (such as Gentamycin™ drug) trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an 30 equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH and the like, suitably are those previously used with the host cell selected for cloning or expression, as the case may be, and will be apparent to the ordinary artisan.

The host cells referred to in this disclosure encompass cells in *in vitro* cell culture as well as cells 35 that are within a host animal or plant.

It is further envisioned that the TRAF polypeptides of this invention may be produced by homologous recombination, or with recombinant production methods utilizing control elements introduced into cells already containing DNA encoding the particular TRAF.

F. Detecting Gene Amplification/Expression

5 Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA **77**, 5201-5205 (1980)]; dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Various labels may be employed, most commonly radioisotopes, particularly ^{32}P . However, other techniques may also be employed, such as using
10 biotin-modified nucleotides for introduction into a polynucleotide. The biotin then serves as a site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to the surface.
15 so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. With immunohistochemical staining techniques, a cell sample is prepared,
20 typically by dehydration and fixation, followed by reaction with labeled antibodies specific for the gene product coupled, where the labels are usually visually detectable, such as enzymatic labels, fluorescent labels, luminescent labels, and the like. A particularly sensitive staining technique suitable for use in the present invention is described by Hse *et al.*, Am. J. Clin. Pharm. **75**, 734-738 (1980).

Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either
25 monoclonal or polyclonal, and may be prepared in any animal. Conveniently, the antibodies may be prepared against a native TRAF polypeptide, or against a synthetic peptide based on the DNA sequence provided herein as described further hereinbelow.

G. Purification of the TRAF polypeptides

The TRAF polypeptide is typically recovered from host cell lysates.

30 When the TRAF polypeptide is expressed in a recombinant cell other than one of human origin, the TRAF is completely free of proteins or polypeptides of human origin. However, it is necessary to purify the TRAF protein from recombinant cell proteins or polypeptides to obtain preparations that are substantially homogenous as to the TRAF. As a first step, the culture medium or lysate is centrifuged to remove particulate cell debris. The membrane and soluble protein fractions are then separated. The TRAF protein
35 may then be purified from the soluble protein fraction. The following procedures are exemplary of suitable purification procedures: fractionation on immunoaffinity or ion-exchange columns; ethanol precipitation;

reverse phase HPLC; chromatography on silica or on a cation exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; and protein A Sepharose columns to remove contaminants such as IgG. Specific purification procedures have been described hereinabove.

- 5 TRAF functional derivatives in which residues have been deleted, inserted and/or substituted are recovered in the same fashion as the native receptor chains, taking into account of any substantial changes in properties occasioned by the alteration. For example, fusion of the TRAF protein with another protein or polypeptide, e.g. a bacterial or viral antigen, facilitates purification: an immunoaffinity column containing antibody to the antigen can be used to absorb the fusion. Immunoaffinity columns such as a rabbit polyclonal
10 anti-TRAF column can be employed to absorb TRAF variant by binding to at least one remaining immune epitope. A protease inhibitor, such as phenyl methyl sulfonyl fluoride (PMSF) also may be useful to inhibit proteolytic degradation during purification, and antibiotics may be included to prevent the growth of adventitious contaminants. The TRAF proteins of the present invention are conveniently purified by affinity chromatography, based upon their ability to specifically associate with the intracellular domain of a TNF-R2.
- 15 One skilled in the art will appreciate that purification methods suitable for native TRAF may require modification to account for changes in the character of a native TRAF or its variants upon expression in recombinant cell culture.

H. Covalent Modifications of TRAF Polypeptides

- Covalent modifications of TRAF are included within the scope herein. Such modifications are
20 traditionally introduced by reacting targeted amino acid residues of the TRAF with an organic derivatizing agent that is capable of reacting with selected sides or terminal residues, or by harnessing mechanisms of post-translational modifications that function in selected recombinant host cells. The resultant covalent derivatives are useful in programs directed at identifying residues important for biological activity, for immunoassays of the TRAF, or for the preparation of anti-TRAF receptor antibodies for immunoaffinity purification of the
25 recombinant. For example, complete inactivation of the biological activity of the protein after reaction with ninhydrin would suggest that at least one arginyl or lysyl residue is critical for its activity, whereafter the individual residues which were modified under the conditions selected are identified by isolation of a peptide fragment containing the modified amino acid residue. Such modifications are within the ordinary skill in the art and are performed without undue experimentation.

- 30 Cysteiny l residues most commonly are reacted with α -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteiny l residues also are derivatized by reaction with bromotrifluoroacetone, α -bromo- β -(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

- 35 Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1M sodium cacodylate at pH 6.0.

Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing α -amino-containing residues include imidoesters such as methyl picolinimide; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pK_a of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ^{125}I or ^{131}I to prepare labeled proteins for use in radioimmunoassay.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ($R'-N=C=N-R'$) such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy and glutaminyl residues by reaction with ammonium ions.

Glutaminyl and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl, threonyl or tyrosyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 [1983]), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group. The molecules may further be covalently linked to nonproteinaceous polymers, e.g. polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in PCT Publication WO90/05534, published 31 May 1990 or U.S. patents 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

Derivatization with bifunctional agents is useful for preparing intramolecular aggregates of the TRAF with polypeptides as well as for cross-linking the TRAF polypeptide to a water insoluble support matrix or surface for use in assays or affinity purification. In addition, a study of interchain cross-links will provide direct information on conformational structure. Commonly used cross-linking agents include 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, homobifunctional imidoesters, and bifunctional maleimides. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimide yield photoactivatable intermediates which are capable of forming cross-links in the presence of light. Alternatively, reactive water insoluble matrices such as cyanogen bromide activated carbohydrates and the systems reactive

substrates described in U.S. Patent Nos. 3,959,642; 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; 4,055,635; and 4,330,440 are employed for protein immobilization and cross-linking.

Certain post-translational modifications are the result of the action of recombinant host cells on the expressed polypeptide. Glutaminyl and aspariginyl residues are frequently post-translationally deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Other post-translational modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl, threonyl or tyrosyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)].

Other derivatives comprise the novel peptides of this invention covalently bonded to a nonproteinaceous polymer. The nonproteinaceous polymer ordinarily is a hydrophilic synthetic polymer, i.e. a polymer not otherwise found in nature. However, polymers which exist in nature and are produced by recombinant or *in vitro* methods are useful, as are polymers which are isolated from nature. Hydrophilic polyvinyl polymers fall within the scope of this invention, e.g. polyvinylalcohol and polyvinylpyrrolidone. Particularly useful are polyvinylalkylene ethers such as polyethylene glycol, polypropylene glycol.

The TRAF polypeptides may be linked to various nonproteinaceous polymers, such as polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The TRAF may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, in colloidal drug delivery systems (e.g. liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th Edition, Osol, A., Ed. (1980).

I. Glycosylation variants of the TRAFs

The native TRAFs are believed to be unglycosylated, however, variants having glycosylation are within the scope herein. For ease, changes in the glycosylation pattern of a native polypeptide are usually made at the DNA level, essentially using the techniques discussed hereinabove with respect to the amino acid sequence variants. Thus, glycosylation signals can be introduced into the DNA sequence of native TRAF polypeptides.

Chemical or enzymatic coupling of glycosides to the TRAF molecules of the molecules of the present invention may also be used to add carbohydrate substituents. These procedures are advantageous in that they do not require production of the polypeptide that is capable of O-linked (or N-linked) glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free hydroxyl groups such as those of cysteine, (d) free sulfhydryl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan or (f) the amide group of glutamine. These methods are described in WO 87/05330 (published 11 September 1987), and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306.

J. Anti-TRAF antibody preparation**(i) Polyclonal antibodies**

5 Polyclonal antibodies to a TRAF molecule generally are raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the TRAF and an adjuvant. It may be useful to conjugate the TRAF or a fragment containing the target amino acid sequence to a protein that is immunogenic in the species to be immunized, e.g. keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glytaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups.

10 Animals are immunized against the immunogenic conjugates or derivatives by combining 1 mg or 1 μg of conjugate (for rabbits or mice, respectively) with 3 volumes of Freud's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of conjugate in Freud's complete adjuvant by subcutaneous injection at multiple sites. 7 to 14 days later the animals are bled and the serum is assayed for anti-TRAF antibody titer. Animals are 15 boosted until the titer plateaus. Preferably, the animal boosted with the conjugate of the same TRAF, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are used to enhance the immune response.

(ii) Monoclonal antibodies

20 Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

For example, the anti-TRAF monoclonal antibodies of the invention may be made using the 25 hybridoma method first described by Kohler & Milstein, *Nature* 256:495 (1975), or may be made by recombinant DNA methods [Cabilly, *et al.*, U.S. Pat. No. 4,816,567].

In the hybridoma method, a mouse or other appropriate host animal, such as hamster is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized in 30 vitro. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, pp.59-103 (Academic Press, 1986)].

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells.

35 For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, J. Immunol. 133:3001 (1984); Brodeur, *et al.*, Monoclonal Antibody Production Techniques and Applications, pp.51-63 (Marcel Dekker, Inc., New York, 1987)].

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against TRAF. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson & Pollard, Anal. Biochem. 107:220 (1980).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods. Goding, Monoclonal Antibodies: Principles and Practice, pp.59-104 (Academic Press, 1986). Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium or RPMI-1640 medium. In addition, the hybridoma cells may be grown in vivo as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

DNA encoding the monoclonal antibodies of the invention is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences. Morrison, *et al.*, Proc. Nat. Acad. Sci. 81, 6851 (1984), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. In that manner, "chimeric" or "hybrid" antibodies are prepared that have the binding specificity of an anti-TRAF monoclonal antibody herein.

Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody of the invention, or they are substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for a TRAF and another antigen-combining site having specificity for a different antigen.

Chimeric or hybrid antibodies also may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate.

5 For diagnostic applications, the antibodies of the invention typically will be labeled with a detectable moiety. The detectable moiety can be any one which is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , or ^{125}I , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; biotin; radioactive isotopic labels, such as, e.g., ^{125}I , ^{32}P , ^{14}C , or ^3H , or an enzyme, such as alkaline
10 phosphatase, beta-galactosidase or horseradish peroxidase.

Any method known in the art for separately conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter, *et al.*, Nature 144:945 (1962); David, *et al.*, Biochemistry 13:1014 (1974); Pain, *et al.*, J. Immunol. Meth. 40:219 (1981); and Nygren, J. Histochem. and Cytochem. 30:407 (1982).

15 The antibodies of the present invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, Monoclonal Antibodies: A Manual of Techniques, pp.147-158 (CRC Press, Inc., 1987).

Competitive binding assays rely on the ability of a labeled standard (which may be a TRAF polypeptide or an immunologically reactive portion thereof) to compete with the test sample analyte (TRAF)
20 for binding with a limited amount of antibody. The amount of TRAF in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies generally are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte which remain unbound.

25 Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three part complex. David & Greene, U.S. Pat No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured
30 using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

(iii) Humanized antibodies

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized
35 antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and

co-workers [Jones *et al.*, Nature **321**, 522-525 (1986); Riechmann *et al.*, Nature **332**, 323-327 (1988); Verhoeyen *et al.*, Science **239**, 1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (Cabilly, *supra*), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

It is important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three dimensional models of the parental and humanized sequences. Three dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e. the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the consensus and import sequence so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding. For further details see U.S. application Serial No. 07/934,373 filed 21 August 1992, which is a continuation-in-part of application Serial No. 07/715,272 filed 14 June 1991.

Alternatively, it is now possible to produce transgenic animals (e.g. mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, e.g. Jakobovits *et al.*, Proc. Natl. Acad. Sci. USA **90**, 2551-2555 (1993); Jakobovits *et al.*, Nature **362**, 255-258 (1993).

(iv) Bispecific antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for a TRAF, the other one is for any other antigen, and preferably for another receptor or receptor subunit. For example, bispecific antibodies specifically binding two different TRAFs, or a TNF receptor (preferably TNF-R2) and a TRAF, are within the scope of the present invention.

Methods for making bispecific antibodies are known in the art.

Traditionally, the recombinant production of bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities [Millstein and Cuello, Nature **305**, 537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in PCT application publication No. WO 93/08829 (published 13 May 1993), and in Traunecker *et al.*, EMBO **10**, 3655-3659 (1991).

According to a different and more preferred approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, and second and third constant regions of an immunoglobulin heavy chain (CH2 and CH3). It is preferred to have the first heavy chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are cotransfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance. In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in copending application Serial No. 07/931,811 filed 17 August 1992.

For further details of generating bispecific antibodies see, for example, Suresh *et al.*, Methods in Enzymology **121**, 210 (1986).

(v) Heteroconjugate antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Patent No. 4,676,980], and for treatment of HIV infection [PCT application publication Nos. WO 91/00360 and WO 92/200373; EP 03089]. Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Patent No. 4,676,980, along with a number of cross-linking techniques.

L. Use of TRAF molecules

One important application of the TRAF proteins of the present invention is based on their ability to associate, directly or indirectly, with the cytoplasmic domain of TNF-R2, CD40 and/or the Epstein-Barr virus transforming protein LMP1. Accordingly, the TRAF proteins can be used in biochemical assays to identify inhibitors of biological activities mediated by TNF-R2, CD40 and/or LMP1. For example, inhibitors of the interaction of TNF-R2 TRAF proteins, such as TRAF2, can be useful in the treatment of various pathological conditions associated with the expression of TNF, such as endotoxic (septic) shock and rheumatoid arthritis (RA). Similarly, inhibitors of the LMP1:TRAF (e.g., TRAF2) interaction might be used to block pathological conditions mediated by the LMP1 oncogene.

Preferably, these assays are performed in the "two-hybrid" assay format, using, for example, the Matchmaker Two-Hybrid System (Clontech), essentially as illustrated in the Examples hereinbelow. Large-scale production and purification of recombinant Glutathione-S-transferase (GST) fusion proteins comprising the cytoplasmic domains of TNF-R2, CD40 or LMP1 are performed as described in the Examples. For large-scale production of a recombinant TRAF protein, the corresponding TRAF cDNA is expressed in Hi5 insect cells (Invitrogen, San Diego) using the baculovirus expression system (commercially available "BaculoGold" Expression Kit from Pharmingen, San Diego). The recombinant TRAF is tagged with a poly histidine tag to allow purification (see the description of anti-TRAF antibody production in Example 4), and a recognition site for heart muscle kinase (Sigma) for radioactive labeling *in vitro*. The biochemical screening assay is performed in a robotic automated system in which the GST-TNF-R2 or /CD40 or /LMP1 fusion protein is coated into 96 well microtiterplates and the radioactively labeled TRAF protein added in the presence of various compounds. After incubation for one hour, the plates are washed to remove unbound TRAF protein and the captured radioactivity is counted. Inhibitors that prevent the interaction between the TRAF protein and the GST-TNF-R2/CD40/LMP1 fusion protein are identified by decreased captured radioactivity compared with control wells that lack added compounds.

In a particular embodiment, the inhibitors of TRAF/TNF-R2, TRAF/CD40, and/or TRAF/LMP1 interaction are functional derivatives, such as amino acid sequence variants, of the native TRAF. One of such amino acid sequence variants is a protein comprising amino acids 87-501 of the native TRAF2 amino acid sequence: TRAF2(87-501) which is a potent inhibitor of the interaction of native TRAF2 with TNF-R2, CD40 and LMP1 (see Example 6). Based upon their ability to specifically associate with the intracellular domain of TNF-R2, the TRAF molecules of the present invention can be used to purify TNF-R2, which, in turn, is useful in the treatment of various pathological conditions associated with the expression of TNF, such as endotoxic (septic) shock and rheumatoid arthritis (RA), either as a soluble TNF-R2 protein or in the form of an immunoglobulin fusion protein. The dose regimens effective in the treatment of these and other diseases can be determined by routine experimentation. CD40 and LMP1 can be purified in an analogous manner, taking advantage of the ability of TRAF proteins, such as TRAF2, to bind these proteins.

The TRAF molecules of the present invention may additionally be used to generate blocking (antagonist) or agonist anti-TRAF antibodies, which can be used to block or mimic TNF biological activities mediated (exclusively or partially) by TNF-R2, or to purify other TRAF proteins having an epitope to which

the antibodies bind. Generic methods for generating anti-TRAF antibodies have been described hereinabove, and are specifically illustrated in Example 4. Such antibodies may be screened for agonist and antagonist properties in assay systems similar to those described above for the identification of inhibitors of the TRAF/TNF-R2/CD40/LMP1 interaction.

5 Therapeutic formulations comprising antibodies, other polypeptides or small organic molecules identified or purified using the TRAF proteins of the present invention, or comprising TRAF proteins (including functional derivatives) of the present invention, are prepared for storage by mixing the active ingredient having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers [Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)], in the form of
10 lyophilized formulations or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides,
15 disaccharides and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, Pluronic or PEG.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and
20 poly-(methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, *supra*.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution.

25 Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

The route of administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial or intralesional routes, topical administration, or by sustained release systems.

30 Suitable examples of sustained release preparations include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices include polyesters, hydrogels, polylactides [U.S. Patent 3,773,919, EP 58,481], copolymers of L-glutamic acid and gamma ethyl-L-glutamate [U. Sidman *et al.*, Biopolymers 22 (1): 547-556 (1983)], poly (2-hydroxyethyl-methacrylate) [R. Langer, *et al.*, J. Biomed. Mater. Res. 15: 167-277 (1981) and R. Langer, Chem. Tech. 12: 98-105 (1982)], ethylene
35 vinyl acetate [R. Langer *et al.*, *Id.*] or poly-D-(-)-3-hydroxybutyric acid [EP 133,988A]. Sustained release compositions also include liposomes. Liposomes containing a molecule within the scope of the present invention are prepared by methods known *per se*: DE 3,218,121A; Epstein *et al.*, Proc. Natl. Acad. Sci. USA 82: 3688-3692 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA 77: 4030-4034 (1980); EP 52322A; EP

36676A; EP 88046A; EP 143949A; EP 142641A; Japanese patent application 83-118008; U.S. patents 4.485.045 and 4.544.545; and EP 102.324A. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. % cholesterol, the selected proportion being adjusted for the optimal NT-4 therapy.

5 An effective amount of the active molecule will depend, for example, upon the therapeutic objectives, the route of administration, and the condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. A typical daily dosage might range from about 1 µg/kg to up to 100 mg/kg or more, depending on the factors mentioned above. Typically, the clinician will administer a molecule of the present invention until a dosage
10 is reached that provides the required biological effect. The progress of this therapy is easily monitored by conventional assays.

Further details of the invention will be apparent from the following non-limiting examples.

Example 1

Purification of TRAF1

15 A. Cell Culture and Biological Reagents

The murine interleukin 2 (IL-2)-dependent cytotoxic T cell line CT6 [Ranges *et al.* J. Immunol. **142**, 1203-1208 (1989)] was cultured in RPMI 1640 media supplemented with 10-20 units recombinant human IL-2 (Boehringer Mannheim), 10-15% fetal calf serum (Hyclone), 2 mM L-glutamine, 10⁻⁵M β-mercaptoethanol, 100 units of penicillin per ml, and 100 µg of streptomycin per ml (GIBCO/BRL). The human T-cell
20 lymphoma line Jurkat was obtained from the American Type Culture Collection (ATCC; Rockville, MD) and maintained in RPMI 1640 media containing 10% fetal calf serum. The human embryonic kidney cell line 293 (ATCC CRL 1573) and 293 cells overexpressing the hTNF-R2 (293/TNF-R2) were maintained as described [Pennica *et al.*, J. Biol. Chem. **267**, 21172-21178 (1992)]. Recombinant hTNF and recombinant mTNF (specific activity of > 10⁷ units/mg) were provided by the Genentech Manufacturing Group. The rabbit
25 anti-human and anti-murine TNF-R2 antibodies have been described previously [Pennica *et al.*, *supra*; Tartaglia *et al.*, J. Biol. Chem. **267**, 4304-4307 (1991)]. Anti-human TNF-R1 monoclonal antibody 986 (IgG2a isotype) and anti-human TNF-R2 monoclonal antibodies 1036, 1035 and 1038 (IgG2b, IgG2a and IgG2b isotypes, respectively) were produced as described [Pennica *et al.*, Biochemistry **31**, 1134-1141 (1992)].

30 B. Mutational Analysis of the Intracellular Domain of hTNF-R2

It has been shown that the TNF induced proliferation of murine CT6 cells is mediated by the 75 kd TNF receptor [TNF-R2; Tartaglia *et al.*, 1991, *supra*]. In addition, TNF-R2 activates the transcription factor NF-κB [Lenardo & Baltimore, Cell **58**: 227-229 (1989)] and mediates the transcriptional induction of the granulocyte-macrophage colony stimulating factor (GM-CSF) gene [Miyatake *et al.*, EMBO J. **4**, 2561-2568
35 1985]; Stanley *et al.*, EMBO J. **4**, 2569-2573 (1985)] and the A20 zinc finger protein gene [Opipari *et al.*, J. Biol. Chem. **265**, 14705-14708 (1990)] in CT6 cells (Figure 1).

To identify sequences within the intracellular domain of the hTNF-R2 (hTNF-R2icd) that are required for TNF signaling a series of mutant hTNF-R2 expression vectors was generated that encode receptors with truncated intracellular domains. DNA fragments containing C-terminally truncated hTNF-R2icds were amplified from the full length expression vector pRK-TNF-R2 [Tartaglia *et al.*, Cell **73**, 213-216 (1993)] by

5 PCR with *Pfu* DNA polymerase (Stratagene). PCR was run for 20 cycles (45 s at 95°C; 60 s at 55°C; 60 s at 72°C) after an initial step of 6 min at 95°C. A 0.5 kb DNA fragment encoding an intracellular domain which lacks amino acids 424-439 of the wild type hTNF-R2 was amplified using the oligonucleotide primers

5'-CCTTGTGCCTGCAGAGAGAAG-3' (SEQ. ID. NO: 23) and 5'-CTAGGTAACTTTCCGGTGCTCCCCAGCAGGGTCTC-3' (SEQ. ID. NO: 24). The fragment was digested

10 with *Pst*I and *Hind*III, gel purified, and re-cloned into the hTNF-R2 cDNA using the expression vector pRIS [Tartaglia & Goeddel, J. Biol. Chem. **267**, 4304-4307 (1992); hTNF-R2(-16)). Similar mutant hTNF-R2 expression vectors were generated that encode receptors lacking amino acids 403-439 (5'-CTAGGTAACTGGAGAAGGGGACCTGCTCGTCCTT-3' (SEQ. ID NO: 25); hTNF-R2(-37)), amino acids

381-439 (5'-CTAGGTAACTGCTGGCTTGGGAGGAGCACTGTGA-3' (SEQ. ID NO: 26); hTNF-R2(-59)), amino acids 346-439 (5'-CTAGGTAACTGCTCCCGGTGCTGG CCCGGGCCTC-3' (SEQ. ID NO: 27); hTNF-R2(-94)) and amino acids 308-439 (5'-CTAGGTAACTGCACTGGCCGAGCTCTCCAGGGA-3' (SEQ. ID NO: 28); hTNF-R2(-132)). A deletion of amino acids 304-345 of hTNF-R2 was constructed by partial digest of pRK-TNF-R2 with *Sac*I and re-ligation of the vector (hTNF-R2(Δ 304-345)). A deletion of the entire intracellular domain of hTNF-R2 was constructed from pRK-TNF-R2 by replacement of sequences

20 between the *Pst*I site adjacent to the transmembrane domain and the *Cl*aI site with a double-stranded oligonucleotide (5'-GTGATGAGAATTCAT-3' (SEQ. ID NO: 29) and 5'-CGATGAATTCTC ATCACTGCA-3') (SEQ. ID NO: 30) containing an in-frame stop codon immediately following Gln²⁷³ (hTNF-R2(-166)). A mutation converting Ser³⁹³ into Ala was introduced into the hTNF-R2 cDNA by site-directed mutagenesis as described [Tartaglia *et al.*, Cell **74**, 845-853 (1993); hTNF-R2(S393A)]. Verification

25 of correctly modified cDNAs was determined by double-strand sequencing using the Sequenase 2.0 Sequencing Kit (U. S. Biochemical).

The expression vectors encoding the intact and truncated hTNF-R2 were introduced into CT6 cells by electroporation. 5 x 10⁶ cells in 0.3 ml RPMI 1640 media were cotransfected with 0.5 μ g of *Sca*I-digested pRK.neo [Tartaglia & Goeddel, 1992, *supra*] and 20 μ g of *Sca*I-digested hTNF-R2 expression vector using

30 the Bio-Rad Gene Pulser with Capacitance Extender (0.4 cm cuvette, 960 μ F, 250 V). Electroporated cells were resuspended in 50 ml media and after 2 days plated into 96-well microtiter plates by limiting dilution in selective media containing 100 μ g/ml G418 (GIBCO/BRL). After three weeks, individual G418-resistant clones were picked and expanded. Clones that express the hTNF-R2 were identified by FACS analysis as described [Table 1; Pennica *et al.*, J. Biol. Chem., 1992, *supra*].

35 Proliferation of CT6 clones expressing the full length and truncated hTNF-R2 was measured by [³H]thymidine incorporation as described [Tartaglia *et al.*, Proc. Natl. Acad. Sci. USA **88**, 9292-9296 (1991)]. NF- κ B activation was analyzed by electrophoretic mobility shift assay with nuclear extracts prepared from stimulated or unstimulated CT6 cells as described [Schütze *et al.*, Cell **71**, 765-776 (1992)].

Table 1 shows that the transfected hTNF-R2 signals proliferation and NF- κ B activation in CT6 cells. In addition, mutant human receptors which lack the C-terminal 16 amino acids or the internal 42 amino acids 304-345 are still functional in mediating these activities. In contrast, mutant receptors which lack the C-terminal 37 amino acids or contain further C-terminal deletions are defective in these assays. These results indicate that a region of 78 amino acids within the intracellular domain of hTNF-R2 comprising amino acids 346-423 is required for mediating TNF signaling. This region contains a potential protein kinase C phosphorylation site (Ser³⁹³-Pro³⁹⁴-Lys³⁹⁵) which is conserved in the murine TNF-R2. However, a mutant hTNF-R2 containing Ala instead of Ser³⁹³ is biologically functional (Table 1) indicating that this phosphorylation site is not involved in TNF-R2 mediated signaling.

10 C. Identification of Factors that Associate with the Intracellular Domain of hTNF-R2

To identify factors that are associated with the intracellular domain of hTNF-R2 immunoprecipitation of the receptor from lysates of [³⁵S]-labeled transfected CT6 cells was performed. 5 x 10⁶ CT6 cells expressing the wild type hTNF-R2 were washed twice with low glucose Dulbecco's modified Eagle's media without cysteine and methionine and incubated in fresh media for 30 min. The cells were seeded into a 100-mm plate in 5 ml media (without cysteine and methionine) containing [³⁵S]cysteine and [³⁵S]methionine (50 μ Ci of L-[³⁵S]-*in vitro* cell labeling mix/ml; Amersham). The cells were incubated for 4 h at 37°C, stimulated for 10 min with 100 ng/ml hTNF, harvested, washed twice with cold PBS and lysed for 20 min at 4°C in 1 ml of 0.1% NP40 lysis buffer containing 50 mM HEPES pH 7.2, 250 mM NaCl, 10% Glycerol, 2 mM EDTA, 1 mM PMSF, 1 μ g/ml Benzamidine, 1 μ g/ml Aprotinin, 1 μ g/ml Leupeptin. Nuclear and cell debris were removed by centrifugation at 10,000 x g for 10 min at 4°C. The cell lysate was precleared for 1 h at 4°C with 50 μ l Pansorbin (Calbiochem). The lysate was incubated for 8 h at 4°C with 1 μ g of each of the anti-hTNF-R2 monoclonal antibodies 1035 and 1038 (directed against different epitopes of the extracellular domain of the hTNF-R2) that had been preabsorbed with 1 ml of unlabeled lysate from untransfected CT6 cells and collected with 15 μ l of protein A-agarose beads (Oncogene Science). The beads were washed extensively with lysis buffer, resuspended in SDS sample buffer and the supernatant electrophoresed on a 4-12% or 8% Tris/glycine polyacrylamide gel. The gel was fixed, incubated in Amplify (Amersham), dried, and exposed to film at -80°C.

Several bands in the range of 45 to 50-56 kd and one band of approximately 68 kd were specifically coprecipitated with the immunoprecipitated hTNF-R2 in CT6 cells (Figure 2a). The same result was obtained when the hTNF-R2 was immunoprecipitated from unlabeled 293/TNF-R2 cells followed by incubation with labeled lysate from untransfected CT6 cells (Figure 2b). The pattern of bands coprecipitated with hTNF-R2 was identical regardless of whether the lysate was prepared from cells that had been stimulated with hTNF or left unstimulated, indicating that these proteins are constitutively associated with the hTNF-R2. This is similar to results observed for the tyrosine kinase JAK2 which is associated with the intracellular domain of the erythropoietin receptor [Witthuhn *et al.*, *Cell* 74, 227-236 (1993)].

In order to establish a large scale purification procedure for factors that associate with the hTNF-R2icd, the intracellular domain of hTNF-R2 was expressed as a glutathione S-transferase (GST) fusion protein

[Smith & Johnson, 1988, *supra*]. The intracellular domain of hTNF-R2 was amplified from pRK-TNF-R2 by PCR with *Pfu* DNA polymerase as described above using the oligonucleotide primers 5'-GATCGGATCCAAAAGAAGCC CTTGTGCCTGCA-3' (SEQ. ID NO: 31) and 5'-GCCTGGTAACTGGGC-3' (SEQ. ID NO: 32). The amplified 0.55 kb DNA fragment was blunt-ended, digested with *Bam*HI and cloned into *Bam*HI/*Sma*I-digested pGEX-2TK vector (Pharmacia; pGST-hTNF-R2icd). The pGST-hTNF-R2icd plasmid was transformed into a protease deficient strain of *E. coli* K12 carrying the *lac*^R gene on the chromosome (Genentech), an overnight culture diluted 1:10 in fresh LB-medium containing 100 µg/ml carbenicillin and grown at 37°C for 2 h. After induction with 0.1 mM IPTG, cells were grown for 1 h at 37°C, pelleted and washed once with cold PBS. The cells were resuspended in 1/100 culture volume of resuspension buffer containing 20 mM Tris-HCl pH 7.5, 1 M NaCl, 5 mM EDTA, 1mM DTT, 1 mM PMSF, 1 µg/ml Benzamidine, 1 µg/ml Aprotinin, 1 µg/ml Leupeptin. After sonication on ice, insoluble material was removed by centrifugation at 10,000 x g for 15 min at 4°C. Triton X-100 was added to 1% and the cell lysate incubated for 30 min at room temperature on a rotator with 500 µl of a 50% slurry of glutathione-agarose beads (sulphur linkage; Sigma) in PBS per 1 l culture volume. The beads were collected by brief centrifugation at 500 x g and washed extensively with resuspension buffer. An aliquot of the purified GST-hTNF-R2icd fusion protein was analyzed by SDS-PAGE (Figure 3). Concentrations of 5-8 mg fusion protein/ml of swollen beads were obtained routinely.

To prepare a covalently linked GST-hTNF-R2icd fusion protein affinity matrix, the fusion protein was eluted from glutathione-agarose beads by competition with free glutathione using 3 x 30 min washes with 1 bead volume of 250 mM Tris-HCl pH 8.0 containing 50 mM reduced glutathione (Sigma). The eluted fusion protein was dialyzed against 0.1 M HEPES pH 7.2, 150 mM NaCl and covalently coupled to Affigel10/15 (2:1 ratio; Bio-Rad) according to the instructions of the manufacturer. Fusion protein concentrations of up to 10 mg/ml of swollen beads were obtained.

Coprecipitation experiments with GST-hTNF-R2icd fusion protein were performed by incubating 3 µl fusion protein beads with 1 ml of cell lysate prepared from [³⁵S]-labeled CT6 cells as described above. After 8 h at 4°C the fusion protein beads were extensively washed with lysis buffer and analyzed by SDS-PAGE and autoradiography. A pattern of bands was found to specifically coprecipitate with the GST-hTNF-R2icd fusion protein either bound to glutathione-agarose beads or covalently coupled to Affigel10/15 (Figure 4) that was very similar in size to the bands coprecipitating with the immunoprecipitated hTNF-R2 (see Figure 3). This suggests that the GST-hTNF-R2icd fusion protein expressed in *E. coli* does associate with the same intracellular factors as the wild type hTNF-R2 in CT6 cells.

Expression vectors were made that encode GST-hTNF-R2icd fusion proteins with mutant intracellular domains according to the mutational analysis described above. Using the same strategy as for the wild type hTNF-R2icd DNA fragments encoding the mutant -16, -37, -59 and Δ304-345 hTNF-R2 intracellular domains were amplified by PCR and cloned into the pGEX-2TK vector. In addition, a 0.14 kb DNA fragment was amplified using the oligonucleotide primers 5'-GATCGGATCCGGAGACACAGATTCCA GCCCC-3' (SEQ. ID NO: 53) and 5'-GATCGAATTCTTAACCTCTTCGGTGCTCCCCAGCAG-3' (SEQ. ID NO: 54), digested with *Bam*HI and *Eco*RI and cloned into pGEX-2TK. This DNA fragment encodes a peptide of 41 amino acids that

correspond to amino acids 384-424 of the hTNF-R2icd. The fusion proteins were expressed, purified and assayed for coprecipitating proteins as described above.

As shown in Figure 5 the GST-hTNF-R2icd fusion proteins containing the intracellular domains of the functional receptor mutants hTNF-R2(-16) and hTNF-R2(Δ 304-345) coprecipitated the same bands as the fusion protein containing the wild type hTNF-R2icd. In contrast, the GST-hTNF-R2icd fusion proteins which contain the intracellular domains of the inactive mutants hTNF-R2(-37) and hTNF-R2(-59) did not coprecipitate these bands. This correlation between the biological activity of hTNF-R2s with mutant intracellular domains and the coprecipitation results obtained with the corresponding GST-hTNF-R2icd fusion proteins supports the observation that the wild type GST-hTNF-R2icd fusion protein associates with the same intracellular factors as the immunoprecipitated hTNF-R2.

In addition, the GST-hTNF-R2icd(384-424) fusion protein was able to coprecipitate the bands at 45 to 50-56 kd and 68 kd although to a weaker extent than the other fusion proteins (Figure 5). The 41 amino acids of the hTNF-R2icd contained in this GST-fusion protein are comprised within the 78 amino acids region of the hTNF-R2icd that has been identified to be required for mediating TNF signaling in CT6 cells (see above). This suggests that this short region of the hTNF-R2icd is sufficient to mediate the association of potential signaling molecules with the receptor.

Competition coprecipitation experiments were performed in which the hTNF-R2 was immunoprecipitated from unlabeled 293/TNF-R2 cells and then incubated with labeled CT6 cell lysate that had been precleared with 50 μ l of GST-hTNF-R2icd fusion protein beads. Preincubation of the CT6 extracts with GST beads alone or GST-hTNF-R2icd(-37) and GST-hTNF-R2icd(-59) fusion protein beads had no effect on the pattern of proteins coprecipitating with the immunoprecipitated hTNF-R2 (Figure 6). However, if the cell lysate had been precleared with GST-hTNF-R2icd or GST-hTNF-R2icd(-16) fusion protein beads, these proteins did not coprecipitate with the immunoprecipitated hTNF-R2 (Figure 6), indicating that they had been depleted from the labeled CT6 cell extract by the GST-hTNF-R2icd fusion proteins. This result demonstrates that the wild type GST-hTNF-R2icd fusion protein associates with the same intracellular factors as the immunoprecipitated hTNF-R2. Consequently, this GST-fusion protein material can be used for large scale purification of factors that are associated with the intracellular domain of hTNF-R2. Coprecipitation experiments of GST-hTNF-R2icd fusion beads with cell lysate prepared from [35 S]-labeled human Jurkat cells revealed a pattern of coprecipitating proteins very similar in size to the pattern observed with murine CT6 lysates (Figure 7). This suggests that the TNF-R2 associated factors are closely related between the mouse and human species.

To investigate the subcellular localization of the hTNF-R2 associated factors, cytoplasmic and membrane fractions from CT6 cells were prepared essentially as described [Deutscher, Methods in Enzymol. 182: Academic Press, San Diego (1990)]. Briefly, [35 S] labeled CT6 cells were washed once with cold PBS and once with isotonic salt buffer containing 50 mM Tris-HCl pH 7.4, 100 mM NaCl, 1 mM EDTA, 1 mM PMSF, 1 μ g/ml Aprotinin and 1 μ g/ml Leupeptin. Cells were resuspended in 5 ml isotonic salt buffer, and lysed in a glass douncer (Wheaton) with 20 strokes using the 'B' pestle. Large cell debris were removed by centrifugation at 750 x g for 10 min at 4°C and the supernatant subjected to ultracentrifugation at 100,000

x g for 30 min at 4°C. The supernatant which constitutes the cytoplasmic fraction was removed and the pellet resuspended in 50 mM TrisHCl pH 7.4, 1 mM EDTA, 1 mM PMSF, 1 µg/ml Aprotinin, 1 µg/ml Leupeptin. This crude membrane fraction was layered on a cushion of 35% w/v sucrose in PBS and centrifuged at 30,000 x g for 45 min at 4°C. The purified cell membrane fraction at the interface between the sucrose and the buffer phases was removed carefully, concentrated by centrifugation at 100,000 x g for 30 min at 4°C and extracted with 0.1% NP40 lysis buffer for 30 min at 4°C. The cell membrane lysate and the cytoplasmic fraction were used in coprecipitation experiments with GST-hTNF-R2cd fusion protein beads as described above.

The factors associating with the hTNF-R2cd were found to be localized in the cytoplasmic cell fraction (Figure 8). A small amount could also be detected in the purified cell membrane fraction consistent with the observation that these factors are constitutively associated with the hTNF-R2cd (see above).

D. Large scale purification

For large scale purification of hTNF-R2cd associated factors 60 l of CT6 cells (3×10^{10} cells) were harvested and washed twice with cold PBS. All subsequent operations were carried out at 4°C. The cells were lysed by adding 120 ml of 0.1% NP40 lysis buffer containing 100 mM NaCl and rocked gently for 30 min. Insoluble material was removed by centrifugation for 10 min at 10,000 x g. The supernatant was then centrifuged at 100,000 x g for 1 hr and dialyzed against lysis buffer containing 500 mM NaCl. All subsequent purification steps were carried out in lysis buffer containing 500 mM NaCl. The cell lysate was passed through a 15 ml glutathione-agarose GST-hTNF-R2cd(-37) fusion protein preabsorption column. The flow-through was applied to a 0.3 ml Affigel10/15 GST-hTNF-R2cd fusion protein affinity column. For control, the lysate was run through a similar Affigel10/15 GST-hTNF-R2cd(-37) fusion protein affinity column in parallel. After extensive washing, proteins bound to the resins were eluted with five column volumes of ImmunoPure Gentle Ag/Ab Elution Buffer (Pierce) containing, 0.1 M DTT, precipitated with Methanol/Chloroform and resuspended in SDS sample buffer containing 5% SDS. One tenth of the material was separated by SDS-PAGE under reducing conditions and visualized by silver staining (Figure 9). Protein bands that were specifically eluted from the GST-hTNF-R2cd fusion protein affinity column were observed at approximately 45 to 50-56 kd and 68-70 kd.

The remaining purified material was separated by SDS-PAGE, electrophoretically transferred to PVDF sequencing membrane (Millipore) and proteins visualized by staining with R250. The protein band at 45 kd (TNF Receptor Associated Factor 1 or TRAF1) was cut out and subjected to amino acid sequence analysis by automated Edman degradation on an Applied Biosystems sequencer. Since the material proved to be N-terminally blocked, internal sequence information was obtained from individual peptides that were purified by reversed phase capillary HPLC after protease digestion prior to sequence analysis. Two peptides that were obtained from trypsin and lysine C digestion, respectively, had the sequences APMALER and KHAYVK (SEQ ID NOS: 41 and 42).

Example 2

Recombinant production of TRAF1

The following degenerate oligonucleotides were designed based on the sequences of the above peptides: BP50-1 sense, 5'-GCNCCNATGGCNYTNGARC/AG (SEQ. ID. NOs: 33-35); BP50-1 antisense, 5'-CT/GYTCNARNGCCATNGGNGC (SEQ. ID NOs: 36-38); BP50-11 sense, 5'-AARCAYGCNTAY GTNAA (SEQ. ID NO: 39); BP50-11 antisense, 5'-TTNACRTANGCRTGYTT (SEQ. ID NO: 40). 1 µg poly(A)⁺mRNA isolated from CT6 cells was oligo(dT)-primed and reverse transcribed using the cDNA Cycle Kit (Invitrogen) according to the instructions of the manufacturer. First-strand CT6 cDNA was subjected to PCR with combinations of the degenerate oligonucleotides listed above using a Cetus GeneAmp Kit and Perkin-Elmer Thermocycler. The PCR was run for 35 cycles (45 s at 95°C; 60 s at 55°C; 150 s at 72°C) after an initial step of 6 min at 95°C. The PCR products were analyzed by electrophoresis on a 1.6% agarose gel. The PCR reaction obtained with the primer combination BP50-1 sense and BP50-11 antisense Coomassie Brilliant Blue R-250 (Sigma) contained an amplified DNA fragment of approximately 0.75 kb. This fragment was gel-purified, subcloned into pBluescript KS (Stratagene), and sequenced.

A 0.65 kb *Pst*I DNA fragment was isolated from the cloned PCR fragment and labeled with [α -³²P]dCTP using the T7 Quick Prime Kit (Pharmacia). The labeled fragment was used to screen approximately 1×10^6 recombinant phage clones from a CT6 cDNA library that had been constructed in λ gt22a using the Superscript Lambda System For cDNA Synthesis And λ Cloning (GIBCO/BRL) according to the instructions of the manufacturer. Hybridization and washing of the filters were carried out under high-stringency conditions according to standard protocols [Ausubel *et al.*, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, New York 1987]. Four positive clones were plaque-purified by a secondary screen. The cDNA inserts of these phage clones were subcloned into pBluescript KS and sequenced on both strands (Figure 10). The longest of the cDNAs was found to be 2 kb. The other three cDNA clones represented truncated versions of the 2 kb cDNA clone. The 2 kb cDNA clone contained an open reading frame encoding a protein of 409 amino acids (Figure 10). Within the predicted protein were the sequences APMALER (SEQ. ID. NO: 41) and KHAYVK (SEQ. ID. NO: 42), as well as the sequences PGSNLGS (SEQ. ID. NO: 43) and KDDTMFLK (SEQ. ID. NO: 44) which correspond to two other peptide sequences obtained from protein sequence analysis. These results confirm that the isolated 2 kb cDNA clone encodes the purified TRAF1.

To analyze similarities between TRAF1 and other known sequences, the TRAF1 sequence was searched against the Genentech protein database. No obvious similarity of significance between TRAF1 and any other known protein was found, indicating that TRAF1 is a novel molecule.

Example 3

Identification and cloning of TRAF2

To directly isolate genes coding for proteins that associate with the intracellular domain of TNF-R2 the yeast two-hybrid system for the detection of protein-protein interactions (Fields & Song, Nature 340, 245-246 [1989]) was used.

The intracellular domain of hTNF-R2 was amplified from pRK-TNF-R2 by PCR with *Pfu* DNA polymerase as described above using the oligonucleotide primers 5'-TCG ATCGTCGACCAAAAAGAAGCCCTCCTGCCTACAA-3' (SEQ. ID NO: 45) and 5'-CTAGAGATCTCAGG GGTCAAGGCACTTT-3' (SEQ. ID. NO: 46). The amplified 0.55 kb DNA fragment was digested with *Sall* and *BglII*, gel-purified and cloned into the GAL4 DNA-binding domain vector pPC97 (Chevray & Nathans, 1992, supra; pPC97-hTNF-R2icd). Similar constructs were made containing the GAL4 DNA-binding domain fused to the hTNF-R2icd(-16) (5'-CTAGA GATCTGTAACTTTCGGTGCTCCCCAGCAGGGTCTC-3' (SEQ. ID. NO: 47); pPC97-hTNF-R2icd(-16)), the hTNF-R2icd(-37) (5'-CTAGAGATCTGTAACTGGAGAAGGGGACCTGCTCGTCC TT-3' (SEQ. ID. NO: 48); pPC97-hTNF-R2icd(-37)), the hTNF-R2icd(-59) (5'-CTAGAGATCTGTAACTGC TGGCTTGGGAGGAGCACTGTGA-3' (SEQ. ID. NO: 49); pPC97-hTNF-R2icd(-59)) and the intracellular domain of the murine TNF-R2 (5'-TCGATCGTCGACCAAAAAGAAGCCCTCCTGCCT ACAA-3' (SEQ. ID NO: 50) 5'-CTAGAGATCTCAGGGGTCAGGCCACTTT-3' (SEQ. ID. NO: 51); pPC97-mTNF-R2icd).

A plasmid cDNA library in the GAL4 transcriptional activation domain vector pPC86 [Chevray & Nathans, 1992, supra] was constructed from *Sall/NorI*-adapted, double-stranded fetal liver stromal cell line 7-4 cDNA (a gift of B. Bennett and W. Matthews) as described [Chevray & Nathans, 1992, supra]. Plasmid DNA was isolated directly from 2×10^6 transformed *E. coli* DH10B (GIBCO/BRL) colonies.

S. cerevisiae HF7c (Clontech) was sequentially transformed with pPC97-hTNF-R2icd and 250 μ g library plasmid DNA as described in the Matchmaker Two-Hybrid System (Clontech). The final transformation mixture was plated onto 50 150-mm synthetic dextrose agar plates lacking L-tryptophan, L-leucine, L-histidine and containing 20 mM 3-aminotriazole (Sigma). A total of 2×10^6 transformed colonies were plated. After 4 days at 30°C 42 surviving His⁺-colonies were obtained of which 15 were positive in a filter assay for β -galactosidase activity [Breden & Nasmyth, Regulation of the Yeast HO gene, Cold Spring Harbor Symposia on Quantitative Biology 50, 643-650. Cold Spring Harbor Laboratory Press, New York (1985)]. Yeast DNA was prepared [Hoffmann and Winston, Gene 57, 267-272 (1987)], transformed into *E. Coli* DH10B by electroporation and colonies containing the pPC86 library plasmid identified by restriction analysis. 14 out of 15 cDNA inserts had a similar size of approximately 2.1 kb. Restriction analysis with *DdeI* revealed them to be independent cDNA clones derived from the same mRNA species.

Retransformation of three representative cDNA clones into HF7c cells with pPC97 and pPC97-hTNF-R2icd, respectively, confirmed that the encoded GAL4 activation domain fusion proteins do not interact with the GAL4 DNA-binding domain alone but only with the GAL4 DNA-binding hTNF-R2icd fusion protein. The 2.1 kb cDNA insert of one representative clone (pPC86Y17) was sequenced on both strands (Figure 11). In addition, the 5'- and 3'-regions of 6 other independent cDNA clones were sequenced confirming that they were derived from the same mRNA species. All clones were shown to be fused to the GAL4 DNA-binding domain in the same reading frame within 20 nucleotides of each other.

Two additional cDNA clones were isolated from a CT6 λ phage cDNA library (see above) and 5 additional clones from a mouse liver λ phage cDNA library (Clontech) using a [³²P]-labeled 0.5 kb *PstI* DNA fragment from the 5'-region of the pPC86Y17 cDNA insert as hybridization probe. None of these cDNA

inserts extended the 5'-sequence of the pPC86 cDNA inserts. Furthermore, the size of the pPC86 cDNA inserts corresponds closely to the size of the actual message as revealed by northern blot analysis of CT6 mRNA (see below). These findings indicate that the cDNA inserts isolated with the two-hybrid system represent full length clones and that the fusion to the GAL4 DNA-binding domain occurred in a very short 5'-untranslated region in-frame with the initiator ATG at position 30 of the pPC86Y17 cDNA insert (see also below). The cDNA clones contain an open reading frame encoding a protein of 501 amino acids (TNF Receptor Associated Factor 2 or TRAF2; Figure 11).

Amino acid sequence analysis of the biochemically purified 56 kd protein associated with the cytoplasmic domain of TNF-R2 (see Figure 9) identified this protein as TRAF2 and confirmed the TRAF2 sequence as predicted from the cDNA sequence (see Figure 11).

A homology search of the TRAF2 sequence against the Genentech protein database revealed that TRAF2 is a novel protein containing an N-terminal RING finger sequence motif [Freemont *et al.*, Cell 64, 483-484 (1991)]; Haupt *et al.*, Cell 65, 753-763 (1991); Inoue *et al.*, *supra*; Figure 12a]. This sequence motif has been observed in the N-terminal domain of a number of regulatory proteins and is thought to form two zinc-binding finger structures that appear to be involved in protein-DNA interactions [Freemont *et al.*, *supra*; Haupt *et al.*, *supra*; Reddy *et al.*, Trends Biochem. Sci. 17, 344-345 (1992)]. Members of the RING finger family are putative DNA-binding proteins, some of which are implicated in transcriptional regulation, DNA repair, and site-specific recombination (see Figure 12 a). In addition, the RING finger motif and other zinc-binding sequence motifs have been discussed to be involved in protein-protein interactions [Freemont *et al.*, *supra*; Haupt *et al.*, *supra*; Berg, J. Biol. Chem. 265, 6513-6516 (1990)]. The importance of this structural motif in TRAF2 is supported by the finding that all GAL4 DNA-binding domain TRAF2 fusions isolated contain the complete N-terminus of TRAF2 (see above). This suggests that the N-terminal RING finger domain of TRAF2 is involved in the interaction with the intracellular domain of TNF-R2.

In addition, TRAF2 shares sequence similarity with the zinc finger motif of *Xenopus* TFIIIA-type zinc finger proteins [Miller *et al.*, EMBO J. 4, 1609-1614 (1985); Berg, *supra*; Figure 12b]. TFIIIA-like zinc finger motifs have also been observed in the RING finger proteins RAD18 and UVS-2 (Figure 12).

No obvious similarity of significance between the C-terminal domain of TRAF2 and any other known protein was found. A comparison of the sequences of TRAF1 and TRAF2 revealed that they share a high degree of amino acid identity in their C-terminal domains (53% identity over 230 amino acids; Figure 13). Both proteins constitute members of a new family of proteins that contain a novel sequence homology motif, the "TRAF domain". The less conserved N-terminal regions within the TRAF domains of TRAF1 and TRAF2 can potentially form leucine zipper-like structures (Figures 10, 11, 13). The leucine zipper is a α -helical structure originally found in a number of DNA-binding proteins that contain leucines occurring at intervals of every seventh amino acid [Landschulz *et al.*, Science 240, 1759-1764 (1988); Vinson *et al.*, Science 246, 911-916 (1989)]. This structure mediates protein dimerization by intermolecular interaction of the leucine side-chains. Leucine zipper structures have also been predicted for two other RING finger family members, the SS-A/Ro ribonucleoprotein and the gene product of the c-cbl proto-oncogene (see Figure 12).

The N-terminal domains of TRAF1 and TRAF2 are unrelated, especially with regard to the RING finger domain of TRAF2.

Example 4

Functional Analysis of TRAF1 and TRAF2

5 Hydropathy profiles [Kyte & Doolittle, 1982, supra] of TRAF1 and TRAF2 (Figure 14) suggest that they lack signal sequences as well as obvious transmembrane regions and are overall hydrophilic. They are thus likely to represent intracellular proteins which is in accordance with the cytoplasmic localization of TRAF1 as determined experimentally (see above).

 Poly(A)⁺mRNA was prepared from CT6 cells [Badley *et al.*, Current Opinion in Structural Biology
10 3, 11-16, (1988)]. Northern analysis [Sambrook *et al.*, 1989, supra] using a radiolabeled TRAF2 hybridization probe as described above indicated that TRAF2 is expressed as a 2.1 kb message in CT6 cells (Figure 15a). Similarly, TRAF1 is expressed in CT6 cells as a 2 kb message (Figure 15a).

 To examine the tissue distribution of TRAF1 and TRAF2 mRNA, mouse multiple tissue Northern blots (Clontech) were hybridized with radiolabeled TRAF1 and TRAF2 probes according to the instructions
15 of the manufacturer. TRAF2 is expressed constitutively in all mouse tissues examined (heart, brain, spleen, lung, liver, skeletal muscle, kidney and testis; Figure 15b). The highest expression level was observed in spleen. In contrast, TRAF1 displayed a tissue specific expression. TRAF1 mRNA could only be detected in spleen, lung and testis (Figure 15b).

 Cotransformation of pPC86Y17 (pPC86TRAF2) into HF7c cells with the above described GAL4
20 DNA-binding TNF-R2^{icd} fusion constructs showed that TRAF2 interacts with the wild type intracellular domains of both the human and the murine TNF-R2 and with the intracellular domain of the biologically active mutant hTNF-R2(-16) (Table 2). However it does not interact with the GAL4 DNA-binding domain alone nor with the intracellular domains of the biologically inactive mutants hTNF-R2(-37) and hTNF-R2(-59). This is in agreement with the results obtained from coprecipitation experiments with wild type and
25 mutant GST-hTNF-R2^{icd} fusion proteins in CT6 cell extracts (see above).

 An expression vector encoding a GST-TRAF2 fusion protein was constructed. The TRAF2 coding region was amplified from pPC86TRAF2 by PCR with *Pfu* DNA polymerase as described above using the oligonucleotide primers 5'-GATCGGATCCTTGTGGTGTGTGGGGG TTGT (SEQ. ID. NO: 55) and 5'-CCTGGCTGGCCTAATGT (SEQ. ID. NO: 56). The amplified 1.6 kb DNA fragment was blunt-ended using
30 *E. coli* DNA polymerase I, digested with *Bam*HI and cloned into *Bam*HI/*Sma*I-digested pGEX-2TK vector. The GST-TRAF2 fusion protein was expressed in the presence of 1 mM ZnCl₂ and purified as described above. GST and GST-TRAF2 fusion protein beads were incubated with lysates from 293 and 293/TNF-R2 cells, and analyzed by SDS-PAGE and Western blot analysis [Sambrook *et al.*, "Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, N.Y. (1989)] using the ECL detection reagent
35 (Amersham). Primary antibodies directed against the extracellular domains of hTNF-R2 and hTNF-R1 (as a control) were used at a concentration of 0.5 µg/ml and the secondary sheep anti-mouse horseradish peroxidase

conjugate (Amersham) at a dilution of 1:6000. As shown in Figure 16, the GST-TRAF2 fusion protein coprecipitates the hTNF-R2 in 293 cell extracts, thus confirming the results obtained from two hybrid analysis.

To test for possible homo- and heteromeric protein-protein interactions between TRAF1, TRAF2 and the intracellular domain of TNF-R2 the 2.1 kb cDNA insert of pPC86TRAF2 was excised by digestion with *Sall* and *NotI* and cloned into pPC97 (pPC97TRAF2). The TRAF1 coding and 3'-untranslated region was amplified from the full length TRAF1 cDNA clone in pBluescript KS by PCR with *Pfu* DNA polymerase as described above using the T7 sequencing primer (Stratagene) and the oligonucleotide primer 5'-TCGATCGTCGACCGCCTCCAGCTCAGCCCCTGAT-3' (SEQ. ID. NO: 52). The amplified 1.7 kb DNA fragment was digested with *Sall* and *NotI*, gel-purified and cloned into both pPC97 and pPC86 (pPC97TRAF1; pPC86TRAF1).

Cotransformation of pPC86TRAF1 into HF7c cells with the GAL4 DNA-binding TNF-R2 fusion constructs encoding the wild type human and murine intracellular domains indicated that the direct interaction between TRAF1 and the intracellular domain of TNF-R2 is weak (Table 2). However, cotransformation of pPC97TRAF1 and pPC86TRAF2 or pPC97TRAF2 and pPC86TRAF1 revealed that TRAF1 and TRAF2 interact with each other (Table 2) suggesting that a heterodimeric complex of TRAF1 and TRAF2 is associated with the intracellular domain of TNF-R2. Subsequently yeast vectors were constructed in which TRAF2 is expressed directly, i. e. not as a GAL4 fusion protein. pPC97TRAF2 was digested with *HindIII* and *Sall* to release a 0.5 kb DNA fragment encoding the GAL4 DNA-binding domain, end-filled with Klenow enzyme, gel-purified, and re-ligated (pPCTRAF2). In addition, TRAF2 was amplified from pPC86TRAF2 with *Pfu* DNA polymerase as described above using the oligonucleotide primers 5'-GATCGACTCGAGATGCCCAAGAAGAAGCGGAAGGTGGC TGCAGCCAGTGTGACTTCCCCT (SEQ. ID. NO: 57) and 5'-CTCTGGCGAAGAAGTCC (SEQ. ID. NO: 58). The amplified 2.1 kb DNA fragment was digested with *XhoI*, end-filled with Klenow enzyme, digested with *NotI*, gel-purified and cloned into pPC97 that had been digested with *HindIII*, end-filled and digested with *NotI* (pPCTRAF2NLS). This expression vector encodes the simian virus 40 large tumor antigen nuclear localization signal (met-pro-lys-lys-lys-arg-lys-val; compare Chevray & Nathans, 1992) fused to the N-terminus of TRAF2. Transformation of pPCTRAF2NLS but not pPCTRAF2 into HF7c cells harboring the plasmids pPC86TRAF1 and pPC97hTNF-R2icd or pPC97mTNF-R2icd complemented the histidine deficiency of the host cells (Table 3). This result confirms that a heterodimeric complex of TRAF1 and TRAF2 interacts with the intracellular domain of TNF-R2. In this protein complex mainly TRAF2 contacts the receptor directly potentially through interaction of its RING finger domain with the C-terminal region of the intracellular domain comprising amino acids 304-345 of the human TNF-R2 as suggested from mutational analysis and coprecipitation experiments (see above). TRAF1 and TRAF2 can also form homodimeric complexes as shown by cotransformation of pPC97TRAF1 and pPC86TRAF1 or pPC97TRAF2 and pPC86TRAF2 (Table 2). These results suggest that the homologous C-terminal domains of TRAF1 and TRAF2 represent a novel protein dimerization motif. In analogy, the C-terminal domain of the RING finger protein COP1 from *Arabidopsis thaliana* contains a region with homology to the β subunit of trimeric G proteins that has been discussed to be involved in protein-protein recognition [Deng *et al.*, *Cell* 71, 791-801 (1992)].

Based on the tissue specific expression of TRAF1 (see above), the formation of a heteromeric complex between TRAF1 and TRAF2 can only occur in certain mouse tissues such as spleen, lung and testis. This raises the possibility of other TRAF domain proteins as tissue specific dimerization partners for the constitutively expressed TRAF2. Such tissue specific heterocomplexes with potentially different biological activities could determine different TNF responses mediated by TNF-R2 in various tissues.

To generate antibodies directed specifically against TRAF1 and TRAF2 the N-terminal domains of both proteins were expressed in *E. coli* as 6xhis tag fusions using the QIAexpress system (Qiagen). A DNA fragment encoding amino acids 2-181 of TRAF1 was amplified from the full length cDNA clone in pBluescript KS by PCR with *Pfu* DNA polymerase as described above using the oligonucleotide primers 5'-GATCGGATCCGCCTCCAGCTCAGCCCCTGAT (SEQ. ID. NO: 59) and 5'-GATCGGATCCAGCCAGCAGCTTCTCCTTCAC (SEQ. ID. NO: 60). The amplified 0.55 kb DNA fragment was digested with *Bam*HI and cloned into *Bam*HI-digested pQE12 vector (Qiagen). Transformants containing the correct orientation of the DNA insert in the expression vector were determined by restriction analysis (pQETRAF1). Similarly, a DNA fragment encoding amino acids 1-162 of TRAF2 was amplified from pPC86TRAF2 using the oligonucleotide primers 5'-GATCGGATCCTTGTGGTGTGTGGGGGTTGT (SEQ. ID. NO: 61) and 5'-GATCGGATCCGCTCAGG CTC TTTTGGGGCA (SEQ. ID. NO: 62), digested with *Bam*HI and cloned into *Bam*HI-digested pQE12 vector (pQETRAF2).

Plasmids pQETRAF1 and pQETRAF2 were transformed into *E. coli* M15[pREP4] (Qiagen). The cells were grown, induced, harvested, and the 6xhis tag TRAF1 and TRAF2 fusion proteins purified by Ni-NTA affinity chromatography (Qiagen) under denaturing conditions according to the instructions of the manufacturer. The purified TRAF1 and TRAF2 fusion proteins were resolved on a 13 % Tris/glycine polyacrylamide gel, stained with 0.05 % Coomassie Brilliant Blue R-250 solution in water, and the appropriate bands excised. Gel slices containing 100-200 µg TRAF1 or TRAF2 fusion protein were used for the immunization of rabbits.

Table 1. Analysis of CT6 Clones Expressing Human TNF-R2 Mutants

CT6 Clone	hTNF-R2 Expression (mean fluorescence)	[³ H]Thymidine Incorporation (fold stimulation)	NF-κB Activation
neo.26	157	0.9	-
hR2.30	303	3.4	+
5 hR2.31	350	4.7	+
-16.25	464	10.7	+
-16.31	465	5.1	+
-37.4	478	1.1	-
-37.20	439	1.0	-
10 -59.1	276	1.3	-
-59.23	374	1.0	-
-94.5	515	0.7	-
-94.6	477	1.0	-
-132.3	296	1.1	-
15 -132.22	344	1.0	-
-166.10	361	1.0	-
-166.13	318	1.0	-
Δ304-345	469	nd	+
S393A.2	407	2.9	nd
20 S393A.8	531	5.5	nd

Expression vectors encoding the intact and truncated hTNF-R2 were transfected into CT6 cells. The expression levels of wild type or mutant receptors of individual CT6 clones were analyzed by flow cytometry and values are expressed as mean fluorescence. For functional analysis two independent CT6 clones were examined for each hTNF-R2 mutant except hTNF-R2(Δ304-345) which represents a pool of sorted cells.

25 Proliferation was measured by [³H]thymidine incorporation of cells that had been treated for 24 hr with a 1:1000 dilution of anti-hTNF-R2 polyclonal antibody. Values are expressed as fold stimulation compared with cells that had been treated with an irrelevant antibody. Data shown are the means of triplicate determinations. Standard deviations were generally less than 5%. NF-κB activation was analyzed by electrophoretic mobility shift assay with nuclear extracts prepared from cells that had been stimulated for 20 min with a 1:500 dilution

30 of anti-hTNF-R2 polyclonal antibody. A plus sign indicates the induction of NF-κB DNA-binding activity compared with nuclear extracts prepared from cells that had been treated with an irrelevant antibody. All CT6 clones retained the ability to induce proliferation and NF-κB activation through the endogenous murine TNF-R2 (data not shown). nd, not determined.

Table 2. Interaction between TRAF1, TRAF2 and the Intracellular

Domain of TNF-R2

	Transformant		Growth on trp ⁻	Growth on trp ⁻
	DNA-binding domain hybrid	Activation-domain hybrid	leu ⁻ medium	leu ⁻ his ⁻ medium
	GAL4(DB)	GAL4(TA)	+	-
5	GAL4(DB)-hTNF-R2icd	GAL4(TA)	+	-
	GAL4(DB)-mTNF-R2icd	GAL4(TA)	+	-
	GAL4(DB)-hTNF-R2icd(-16)	GAL4(TA)	+	-
	GAL4(DB)-hTNF-R2icd(-37)	GAL4(TA)	+	-
	GAL4(DB)-hTNF-R2icd(-59)	GAL4(TA)	+	-
10	GAL4(DB)	GAL4(TA)-TRAF1	+	-
	GAL4(DB)	GAL4(TA)-TRAF2	+	-
	GAL4(DB)-hTNF-R2icd	GAL4(TA)-TRAF1	+	-/+
	GAL4(DB)-mTNF-R2icd	GAL4(TA)-TRAF1	+	-/+
	GAL4(DB)-hTNF-R2icd	GAL4(TA)-TRAF2	+	+
15	GAL4(DB)-mTNF-R2icd	GAL4(TA)-TRAF2	+	+
	GAL4(DB)-hTNF-R2icd(-16)	GAL4(TA)-TRAF2	+	+
	GAL4(DB)-hTNF-R2icd(-37)	GAL4(TA)-TRAF2	+	-
	GAL4(DB)-hTNF-R2icd(-59)	GAL4(TA)-TRAF2	+	-
	GAL4(DB)-TRAF1	GAL4(TA)	+	-
20	GAL4(DB)-TRAF2	GAL4(TA)	+	-
	GAL4(DB)-TRAF1	GAL4(TA)-TRAF1	+	+
	GAL4(DB)-TRAF1	GAL4(TA)-TRAF2	+	+
	GAL4(DB)-TRAF2	GAL4(TA)-TRAF2	+	+
	GAL4(DB)-TRAF2	GAL4(TA)-TRAF1	+	+
25	HF7c cells were cotransformed with plasmids (see text) encoding various GAL4 DNA-binding domain (DB) and GAL4 transcriptional activation domain (TA) fusion proteins as indicated. Aliquots of the same transformation mixture were plated onto synthetic dextrose plates lacking trp and leu and plates lacking trp, leu, his and containing 20 mM 3-aminotriazole. Plus signs indicate growth of transformed yeast colonies on the respective plates. Very similar numbers of transformants from the same transformation mixture (90-			
30	100%) were obtained on plates lacking trp, leu, his and on plates lacking trp and leu only. Minus/plus signs indicate that the number of transformants growing on plates lacking trp, leu, his was approximately 1-2% of the number of colonies obtained from the same transformation mixture on plates lacking trp and leu only.			

Filter assays for β -galactosidase activity were performed on colonies growing on plates lacking all three amino acids. All colonies developed a blue color (data not shown).

Table 3. Interaction between TRAF1, TRAF2 and the Intracellular Domain of TNF-R2 (continued)

	Transformant			Growth on trp- leu ⁻ his ⁻ medium
	DNA-binding domain hybrid	Activation-domain hybrid	Direct expression	
5	GAL4(DB)	GAL4(TA)-TRAF1		-
	GAL4(DB)	GAL4(TA)-TRAF1	NLS-TRAF2	-
	GAL4(DB)	GAL4(TA)-TRAF1	TRAF2	-
	GAL4(DB)-hTNF-R2icd	GAL4(TA)-TRAF1		-/+*
10	GAL4(DB)-hTNF-R2icd	GAL4(TA)-TRAF1	NLS-TRAF2	+
	GAL4(DB)-hTNF-R2icd	GAL4(TA)-TRAF1	TRAF2	-/+*
	GAL4(DB)-mTNF-R2icd	GAL4(TA)-TRAF1		-/+*
	GAL4(DB)-mTNF-R2icd	GAL4(TA)-TRAF1	NLS-TRAF2	+
	GAL4(DB)-mTNF-R2icd	GAL4(TA)-TRAF1	TRAF2	-/+*
15	HF7c cells were sequentially transformed with plasmids (see text) encoding the indicated GAL4 DNA-binding domain (DB) fusion proteins, the GAL4 transcriptional activation domain (TA) TRAF1 fusion protein, and TRAF2 or TRAF2 fused to the simian virus 40 large tumor antigen nuclear localization signal (NLS). The final transformation mixtures were plated onto synthetic dextrose plates lacking trp, leu, his and containing 20 mM 3-aminotriazole. Plus signs indicate growth of transformed yeast colonies on the respective plates.			
20	*See Table 2.			

Example 5

Structure-function analysis of TRAF2

To identify regions within the TRAF2 protein that are required for homodimerization, heterodimerization with TRAF1 and for interaction with the cytoplasmic domain of TNF-R2 GAL4-TRAF2 fusion protein vectors were constructed which express mutant TRAF2 proteins. A 1.9 kb DNA fragment which encodes amino acids 87-501 of TRAF2 was amplified from pPC86TRAF2 by PCR with *Pfu* DNA polymerase as described in the previous examples, using the oligonucleotide primers 5'-GATCGAGTCGACCAGTAGTTCGGCCTTTCAA GAT (SEQ. ID. NO: 63) and 5'-CTCTGGCGAAGAAGTCC (SEQ. ID NO: 64). Similarly, a 1.4 kb DNA fragment encoding amino acids 264-501 of TRAF2 was amplified from this plasmid using the oligonucleotide primers 5'-

- GATCGAGTCGACCGTGGGGCCAGAGCTACTCCAG (SEQ. ID. NO: 65) and 5'-GTACGTGCGGCCGCCTACCACCTGGTTCAAGGTTTCCTG (SEQ. ID. NO: 66). The amplified DNA fragments were digested with *Sall* and *NorI*, gel-purified and cloned into pPC97 and pPC86 (pPC97TRAF2(87-501); pPC86TRAF2(87-501); pPC97TRAF2(264-501); pPC86TRAF2(264-501)).
- 5 Cotransformation experiments into HF7c cells as listed in table 4 indicate that the RING finger domain of TRAF2 is not required for interaction with the cytoplasmic domain of TNF-R2 since the mutant TRAF2 protein in which the RING finger domain (amino acids 1-86) was removed was still able to associate with the cytoplasmic domain of TNF-R2. Also, this mutant TRAF2 protein could still associate with both TRAF1 and wild-type TRAF2. The same results were obtained for the mutant TRAF2 protein which only comprises
- 10 the TRAF domain and a few additional amino acids (amino acids 264-501; Table 4). These results demonstrate that the TRAF domain of TRAF2 is sufficient to mediate homo- and heterodimerization of TRAF1 and TRAF2 as well as for interaction of TRAF2 with the cytoplasmic domain of TNF-R2.

Table 4. Interaction between TRAF1, TRAF2 and the Intracellular Domain of TNF-R2 (continued)

	Transformant		Growth on trp-leu- medium	Growth on trp-leu- his-medium
	DNA-binding domain hybrid	Activation-domain hybrid		
15	GAL4(DB)	GAL4(TA)-TRAF2(87-501)	+	-
	GAL4(DB)-hTNF-R2icd	GAL4(TA)-TRAF2(87-501)	+	+
	GAL4(DB)-mTNF-R2icd	GAL4(TA)-TRAF2(87-501)	+	+
	GAL4(DB)-hTNF-R2icd(-16)	GAL4(TA)-TRAF2(87-501)	+	+
20	GAL4(DB)-hTNF-R2icd(-37)	GAL4(TA)-TRAF2(87-501)	+	-
	GAL4(DB)-TRAF1	GAL4(TA)-TRAF2(87-501)	+	+
	GAL4(DB)-TRAF2	GAL4(TA)-TRAF2(87-501)	+	+
	GAL4(DB)-TRAF2(87-501)	GAL4(TA)-TRAF2(87-501)	+	+
	GAL4(DB)	GAL4(TA)-TRAF2(264-501)	+	-
25	GAL4(DB)-hTNF-R2icd	GAL4(TA)-TRAF2(264-501)	+	+
	GAL4(DB)-mTNF-R2icd	GAL4(TA)-TRAF2(264-501)	+	+
	GAL4(DB)-hTNF-R2icd(-16)	GAL4(TA)-TRAF2(264-501)	+	+

	GAL4(DB)-hTNF-R2icd(-37)	GAL4(TA)-TRAF2(264-501)	+	-
	GAL4(DB)-TRAF1	GAL4(TA)-TRAF2(264-501)	+	+
	GAL4(DB)-TRAF2	GAL4(TA)-TRAF2(264-501)	+	-
	GAL4(DB)-TRAF2(87-501)	GAL4(TA)-TRAF2(264-501)	+	+
5	GAL4(DB)-TRAF2(264-501)	GAL4(TA)-TRAF2(264-501)	+	+

The procedures described in Example 4 and after Table 2 above were used to generate the data set forth in the present table.

While the invention has necessarily been described in conjunction with preferred embodiments and specific working examples, one of ordinary skill, after reading the foregoing specification, will be able to effect various changes, substitutions or equivalents, and alterations to the subject matter set forth herein, without departing from the spirit and scope herein. Hence, the invention can be practiced in ways other than those specifically described herein. All such modifications are intended to be within the scope of the present invention.

All references cited herein and the references cited therein are hereby expressly incorporated by reference.

Example 6

The Functional Role of TRAFs

A. The Interaction of TRAFs

TRAF2 associates directly with the cytoplasmic domain of TNF-R2, while the interaction between TRAF1 and TNF-R2 is mediated via TRAF2 [Rothe *et al.*, Cell 78: 681 (1994)]. In addition to forming stable heteromeric complexes, both TRAF1 and TRAF2 can form homodimers. We used the yeast two-hybrid system to determine whether TRAF3 [originally termed CD40bp, CRAF, or LAP1, Hu *et al.*, J. Biol. Chem. 269: 30069 (1994); Cheng *et al.*, Science 267: 1494 (1995); and Mosialos *et al.*, Cell 80: 389 (1995)] interacts with TRAF1 or TRAF2. Expression vectors for TRAFs and the cytoplasmic domains of members of the TNF receptor superfamily in the yeast two-hybrid system have been reported previously [Rothe *et al.*, Cell 78: 681 (1994); Hu *et al.*, J. Biol. Chem. 269: 30069 (1994)]. Two-hybrid interaction analysis between bait- and prey-encoded fusion proteins was performed as described [Rothe *et al.*, *supra*. Hu *et al.*, J. Biol. Chem. 269: 30069 (1994)]. Although TRAF3 self-associated it did not form heterotypic complexes with either TRAF1 or TRAF2 (Table 5). This finding suggests the existence of oligomerization interfaces within the TRAF family members that specify permissible homo- and/or heteromeric combinations.

B. The interaction of TRAFs with other members of the TNF receptor superfamily

We next tested the interaction of each TRAF with other members of the TNF receptor superfamily. None of the three TRAFs associated with the cytoplasmic domain of either TNF-R1 or the Fas antigen (Table 5). Also, TRAF1 did not interact with CD40, nor did TRAF3 interact with TNF-R2. However, TRAF2 was
5 found to associate strongly with the cytoplasmic domains of both CD40 and TNF-R2 (Table 5).

Both TNF and the CD40 ligand can signal NF- κ B activation. Induction of NF- κ B activity by TNF has been demonstrated in numerous cell types including human embryonic kidney 293 cells [Hsu *et al.*, Cell, in press (1995)] (Fig. 17). As revealed by stimulation with receptor-specific agonistic antibodies, TNF-induced NF- κ B activation in 293 cells is predominantly mediated by TNF-R1 (Fig. 17) possibly due to the
10 low level of endogenous TNF-R2 (see below) [Pennica *et al.*, J. Biol. Chem. 267: 21172 (1992)]. To examine a functional role for TRAFs in NF- κ B activation, 293 cells were transiently transfected with TRAF expression vectors. After 24h, nuclear extracts were prepared and assayed for NF- κ B activity by electrophoretic mobility shift assay.

Expression vectors for TRAF1, TRAF2 and TRAF2(87-501) contained the respective coding regions
15 under the transcriptional control of the cytomegalovirus (CMV) immediate early promotor-enhancer in pRK [Schall *et al.*, Cell 61: 361 (1990)]. Expression of TRAF proteins in transiently transfected 293 cells was confirmed by Western blot analysis with polyclonal antibodies directed against TRAF1 and TRAF2, respectively (Rothe and Goeddel, unpublished results). Expression vectors for TRAF3, CD40 [Hu *et al.*, J. Biol. Chem. 269: 30069 (1994)] and TNF-R2 [Tartaglia *et al.*, Cell 73: 213 (1993)] have been described
20 previously.

293 cells were transiently transfected with TRAF expression vectors. After 24 h nuclear extracts were prepared, and NF- κ B activation was analyzed by electrophoretic mobility shift assay [Schütze *et al. supra*]. The composition of the activated NF- κ B complex was examined by supershift analysis with antibodies that recognize specific NF- κ B subunits (Santa Cruz Biotechnology). Expression of TRAF1 or
25 TRAF3 did not lead to induction of NF- κ B DNA-binding activity. In contrast, TRAF2-expressing 293 cells contained a significant amount of activated NF- κ B (Fig. 17). The major component of the active NF- κ B complex appeared to be the p65:p50 heterodimer as indicated by supershift experiments with antibodies directed against specific NF- κ B subunits (Fig. 17).

C. Activation of NF- κ B dependent reporter genes by TRAF2 expression

30 To determine if TRAF2 expression might activate an NF- κ B dependent reporter gene, an E-selectin-luciferase reporter construct [Schindler and Baichwal, supra] was cotransfected with the various TRAF expression vectors into 293 cells. For cotransfection experiments 3×10^5 293 cells were transfected with 0.5 μ g E-selectin-luciferase reporter gene plasmid [Schindler and Baichwal, Mol. Cell. Biol. 14: 5820 (1994)] and 1 μ g for each expression construct. DNA concentrations were kept constant by supplementation with
35 pRK. Cells were lysed 24 h after transfection and reporter gene activity determined using the Luciferase Assay System (Promega). A pRSV- β gal vector (0.5 μ g) was used for normalizing transfection efficiency. TRAF2 expression potently activated the reporter gene, whereas expression of TRAF1 or TRAF3 had no

effect (Fig 18a). In all cases, reporter gene activity could be (co)induced through the TNF-R1 pathway by the addition of TNF (Fig. 18a). The observed reporter gene induction was dependent on NF- κ B activation since TRAF2 expression failed to activate a control reporter construct in which the NF- κ B sites in the E-selectin promoter were mutated. These results demonstrate that overexpression of TRAF2 alone, but not of TRAF1 or TRAF3, is sufficient to induce NF- κ B activation in 293 cells.

TRAF2 and TRAF3 contain N-terminal RING finger motifs that are not required for interaction with TNF-R2 and CD40 [Rothe *et al.*, *supra*; Hu *et al.*, *supra*; Cheng *et al.*, *Science* 267: 1494 (1995); and Mosialos *et al.*, *Cell* 80: 389 (1995)]. A truncated variant of TRAF2 which lacks the N-terminal 86 amino acids comprising the RING finger domain (TRAF2(87-501)) retains its ability to associate with the cytoplasmic domains of TNF-R2 and CD40 as well as with TRAF1 and wild-type TRAF2 (Table 5) (Rothe *et al.*, *supra*). Interestingly, this otherwise functional TRAF2(87-501) protein is completely defective in inducing NF- κ B dependent reporter gene activity in 293 cells (Fig. 18a). Thus, the RING finger domain of TRAF2 appears to be required for NF- κ B activation.

To confirm the results obtained in 293 cells, similar transient transfection experiments were performed in the murine cytotoxic T cell line CT6. CT6 cells do not express detectable levels of TNF-R1 [Lewis *et al.*, *Proc. Natl. Acad. Sci. USA* 88: 2830 (1991)], and TNF-induced NF- κ B activation in this cell line is mediated by TNF-R2 [Rothe *et al.*, *supra*]. As observed with 293 cells, expression of wild-type TRAF2 potently induced the E-selectin-luciferase reporter gene in CT6 cells, whereas TRAF1, TRAF3, and TRAF2(87-501) did not (Fig. 18b). Transient transfection of CT6 cells was performed using the DEAE-dextran method [F.M. Ausubel *et al.*, *Current Protocols in Molecular Biology* (Green Publishing Associates/Wiley & Sons, Inc., New York, (1994)]. 10⁷ CT6 cells were transfected with a total of 10 μ g of plasmid DNA in 250 μ g/ml DEAE-dextran for 90 min. Reporter gene activity was assayed 40 h after transfection. Furthermore, expression of mutant TRAF2(87-501), but not wild-type TRAF1, -2 or -3, suppressed the NF- κ B dependent reporter gene activity induced by TNF stimulation of CT6 cells (Fig. 18b). Thus, TRAF2(87-501) acts as a dominant negative inhibitor of TNF-R2 mediated NF- κ B activation in CT6 cells.

In contrast to CT6 cells, TNF-induced NF- κ B activation in 293 cells is mediated by TNF-R1 (see above). Overexpression of mutant TRAF2(87-501) in 293 cells reduced TNF-induced E-selectin-luciferase reporter gene activity only slightly (Fig. 18a) indicating that TNF-R1 signaling is not blocked by the dominant negative TRAF2 mutation. To test the effect of TRAFs on TNF-R2 signaling in 293 cells, TNF-R2 was transiently coexpressed with wild-type or mutant TRAFs. Overexpression of TNF-R2 alone was sufficient to initiate NF- κ B dependent reporter gene activity even without concomitant TNF stimulation (Fig. 19a). Similarly, TNF-R1 overexpression in the absence of exogenous TNF is known to induce apoptosis and NF- κ B activation in both HeLa cells and 293 cells [Boldin *et al.*, *J. Biol. Chem.* 270: 387 (1995); Hsu *et al.*, *supra*]. Coexpression of mutant TRAF2(87-501) completely abolished the reporter gene activation induced by TNF-R2 (Fig. 19a). Thus, as in CT6 cells, TRAF2(87-501) exerts a dominant negative effect on TNF-R2 signaling in 293 cells. However, NF- κ B dependent reporter gene activity in 293 cells that coexpressed hTNF-R2 and TRAF2(87-501) could still be induced through the TNF-R1 pathway by the

addition of TNF (Fig. 19a). These observations demonstrate that while the TRAF2 signaling pathway is essential for TNF-R2 mediated responses, it is not involved in TNF-R1 signaling.

D. TRAF2 mediates NF- κ B Activation by CD40

We next examined whether TRAF2 also mediates NF- κ B activation by CD40 since it associates with both TNF-R2 and CD40 (see above). As observed for TNF-R2, overexpression of CD40 in 293 cells is sufficient to induce NF- κ B activation in the absence of ligand (Fig. 19b). Coexpression of CD40 and mutant TRAF2(87-501) completely inhibited CD40-induced reporter gene activity (Fig. 19b), demonstrating that TRAF2(87-501) exerts a dominant negative effect on CD40 signaling as well as TNFR2 signaling. These results indicate that TRAF2 is a common mediator of NF- κ B activation for both TNF-R2 and CD40.

Coexpression of wild-type TRAF2 and TNF-R2 or CD40 in 293 cells did not result in a substantially stronger E-selectin-luciferase reporter gene activity than expression of TRAF2 alone (Fig. 18a, 19a, b), suggesting that downstream elements in the signaling cascade are limiting in 293 cells. Our results also indicate that TRAF1:TRAF2 heterodimers may be functional in signaling NF- κ B activation. Overexpression of TRAF1, which would be expected to convert any endogenous TRAF2 homodimers into TRAF1:TRAF2 heterodimers, did not inhibit TNF-R2 mediated reporter gene activity in either CT6 or 293 cells (Fig. 18b, 19a), although coexpression of TRAF1 and CD40 resulted in a weak reduction of CD40-induced NF- κ B activation (Fig. 19b). Rather, a slight but reproducible increase of reporter gene activity was observed when TNF-R2 and TRAF1 were coexpressed in 293 cells (Fig. 19a).

A surprising result is that wild-type TRAF3 suppressed CD40-induced NF- κ B dependent reporter gene activity in 293 cells (Fig. 19b). Thus, TRAF2 acts as positive signal transducer mediating CD40-induced NF- κ B activation, whereas TRAF3 antagonizes this activity, presumably by binding to overlapping TRAF docking sites on CD40. TRAF3 also inhibited TNF-R2 induced NF- κ B activation in 293 cells (Fig. 19a), although this effect was not observed in CT6 cells (Fig. 18b). While no direct interaction of TRAF3 with TRAF1, TRAF2 or TNF-R2 was detected by two-hybrid analysis (see above), the dominant negative effect of TRAF3 in 293 cells might imply that TRAF3 interacts weakly with one of these components. Inhibition of TNF-R2 mediated NF- κ B activation could then be explained by the high level of overexpression of TRAF3 in 293 cells relative to CT6 cells. It is also possible that TRAF3 participates in signaling pathways distinct from the TRAF2 mediated NF- κ B activation pathway. By analogy with TRAF2, the initiation of such a pathway may involve the RING finger domain of TRAF3. Cheng et al. [Cheng *et al.*, *supra*] recently observed that a truncated TRAF3 protein lacking the N-terminal half of the molecule suppressed CD40 mediated induction of CD23. Whereas Cheng et al. [Cheng *et al.*, *supra*] did not report the effect of native TRAF3, our results could indicate that induction of CD23 expression by CD40 may be mediated by TRAF2 and inhibited by native TRAF3.

E. Interaction of TRAF2 with LMP1

It has been reported that the C-terminal cytoplasmic domain of the Epstein-Barr virus transforming protein LMP1 interacts with TRAF1 and TRAF3 [Mosialos *et al.*, *Cell* 80, 389 (1995)]. The interaction between TRAF1 and LMP1 was found to be indirect and hypothesized to be mediated by a yet unknown

TRAF protein. We used the yeast two-hybrid system to examine if TRAF2 interacts with LMP1. As shown in Table 6, TRAF2, TRAF2(87-501) and TRAF3 associate strongly with the C-terminal cytoplasmic domain of LMP1. Based on these findings, TRAF2 is believed to mediate the TRAF1-LMP1 interaction. LMP1 is a dominant oncogene that has multiple downstream effects on cell growth and gene expression, at least some of which require NF- κ B activation [Laherty *et al.*, *J. Biol. Chem.* 267: 24157 (1992); Rowe *et al.*, *J. Virol.* 68: 5602 (1994)]. Our comparative analysis of all three known TRAFs suggests that TRAF2 initiates NF- κ B activation as a common signal transducer for TNF-R2, CD40 and, potentially, LMP1.

Discussion

An important activity initiated by several members of the TNF receptor superfamily is activation of the transcription factor NF- κ B. The results reported here demonstrate that overexpression of TRAF2, but not TRAF1 or TRAF3, is sufficient to induce NF- κ B activation. In addition, a truncated TRAF2 protein that lacks the N-terminal RING finger domain suppresses TNF-R2 and CD40 mediated NF- κ B activation by acting as a dominant negative mutation. We therefore conclude that TRAF2 is a common mediator of TNF-R2 and CD40 signaling that is crucially involved in the activation of NF- κ B. Conceptually, this signaling cascade is activated by TRAF2 aggregation, which can be triggered by either receptor clustering or TRAF2 overexpression. We have also shown that TRAF2 is likely to be a signal transducer for LMP1.

The findings outlined in this study demonstrate that the involvement of TRAF2 in TNF-mediated NF- κ B activation is restricted to TNF-R2 signaling and is not required for NF- κ B activation by TNF-R1. This observation is supported by the recent identification of a novel protein, TRADD, which is unrelated to the TRAF family and which mediates NF- κ B activation by TNF-R1 [Hsu *et al.*, *supra*]. The identification of two distinct signal transducers, TRAF2 and TRADD, provides molecular evidence for earlier indications that activation of NF- κ B can be achieved through multiple independent signaling pathways [Thanos and Maniatis, *ibid.* 80: 529 (1995)]. It is noteworthy that a mutant TRAF2 protein lacking its N-terminal RING finger domain retains the ability to interact with TNF-R2, CD40, TRAF1, and TRAF2 but is completely defective in signaling NF- κ B activation. This finding suggests that the RING finger domain of TRAF2 is involved in NF- κ B activation and represents a potential protein-protein interaction domain that connects TRAF2 with subsequent steps in the signaling cascade.

Table 5. Interaction between TRAFs and members of the TNF receptor superfamily. Yeast Y190 cells were cotransformed with expression vectors encoding the indicated Gal4 DNA-binding domain and Gal4 transcriptional activation domain fusion proteins. Each transformation mixture was plated on a synthetic dextrose plate lacking tryptophan and leucine. Filter assays for β -galactosidase activity were performed to detect interaction between fusion proteins. Plus signs indicate strong color development within 1 h of the assay. Minus signs indicate no development of color within 24 h. Control transformations with empty Gal4 vectors were negative and are not listed.

	DNA-Binding		Activation Domain Hybrid		
	Domain Hybrid	TRAF1	TRAF2	TRAF2(87-501)	TRAF3
5		TRAF1	+	+	
	+	-			
		TRAF2	+	+	
	+	-			
		TRAF3	-	-	-
10		+			
		TNF-R2	-	+	
	+	-			
		CD40	-	+	
	+	+			
15		TNF-R1	-	-	-
		-			
		Fas	-	-	-
		-			

- 20 Table 6. Interaction between TRAFs and the cytoplasmic domain of LMP1. Yeast Y190 cells were cotransformed with expression vectors encoding the indicated Gal4 DNA-binding domain and Gal4 transcriptional activation domain fusion proteins (14). The Gal4 DNA-binding domain-LMP1 construct contained the C-terminal 176 amino acids of LMP1 (S. Fennewald, V. van Santen, E. Kieff. Nucleotide sequence of an mRNA transcribed in latent growth-transforming virus infection indicates that it may encode
- 25 a membrane protein. J. Virol. 51, 411-419, 1984). Each transformation mixture was plated on a synthetic dextrose plate lacking tryptophan and leucine. Filter assays for b-galactosidase activity were performed to detect interaction between fusion proteins. Plus signs indicate strong color development within 1 h of the assay. Minus signs indicate no development of color within 24 h. Control transformations with empty Gal4 vectors were negative and are not listed.

DNA-Binding	Activation Domain Hybrid			
	TRAF1	TRAF2	TRAF2(87-501)	TRAF3
Domain Hybrid				
LMP1		+	+	+

- 5 All documents cited in this application along with the references cited therein are hereby expressly incorporated by reference.

The following deposits were made on 10 February 1995 at the American Type Culture Collection, Rockville, MD, U.S.A., under the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

- | | |
|---------------------------|-------------|
| 10 Designation: | ATCC Number |
| TRAF2 plasmid pPC86-TRAF2 | 97053 |
| TRAF1 plasmid pBst-TRAF1 | 97054 |

- The making of these deposits is by no means an admission that deposits are required to enable the present invention. On the contrary, the invention is fully enabled without reference to these
- 15 deposits.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Genentech, Inc.

5 (ii) TITLE OF INVENTION: Tumor Necrosis Factor Receptor-Associated Factors

(iii) NUMBER OF SEQUENCES: 66

(iv) CORRESPONDENCE ADDRESS:

10 (A) ADDRESSEE: Genentech, Inc.
(B) STREET: 460 Point San Bruno Blvd
(C) CITY: South San Francisco
(D) STATE: California
(E) COUNTRY: USA
(F) ZIP: 94080

(v) COMPUTER READABLE FORM:

15 (A) MEDIUM TYPE: 3.5 inch, 1.44 Mb disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: patin (Genentech)

(vi) CURRENT APPLICATION DATA:

20 (A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

25 (A) APPLICATION NUMBER: 08/250858
(B) FILING DATE: 27-MAY-1994

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 08/331394
(B) FILING DATE: 28-OCT-1994

(viii) ATTORNEY/AGENT INFORMATION:

30 (A) NAME: Dreger, Ginger R.
(B) REGISTRATION NUMBER: 33,055
(C) REFERENCE/DOCKET NUMBER: 897P2PCT

(ix) TELECOMMUNICATION INFORMATION:

35 (A) TELEPHONE: 415/225-3216
(B) TELEFAX: 415/952-9881
(C) TELEX: 910/371-7168

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 2088 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCCAGCCCCG TTCTCTGCCC CAAGGACGCT ACCGCCCAAT GCGAGCAGAA 50

45 GGCGGCGCAC AGATACAGAA AGTGAGGCTC AGACATATTG AAGACCGTGT 100

GACATAGGGT AGCCAAATGA CAGTGTGAGA AAGTGACATT TACTCAAGGC 150

CACCCAGATA TCCTGGAGGA CCCAGAACCC TGGAGATTCC CATCAGAAAG 200

ACCTTCTGGC CACCTGAAAC CCCAAGATGG CCTCCAGCTC AGCCCCTGAT 250
 GAAAACGAGT TTCAATTTGG TTGCCCCCT GCTCCCTGCC AGGACCCATC 300
 GGAGCCCAGA GTTCTCTGCT GCACAGCCTG TCTCTCTGAG AACCTGAGAG 350
 ATGATGAGGA TCGGATCTGT CCTAAATGCA GAGCAGACAA CCTCCATCCT 400
 5 GTGAGCCCAG GAAGCCCTCT GACTCAGGAG AAGGTTCACT CTGATGTAGC 450
 TGAGGCTGAA ATCATGTGCC CCTTTGCAGG TGTGGCTGT TCCTTCAAGG 500
 GGAGCCCACA ATCCATGCAG GAGCATGAGG CTACCTCCCA GTCCTCCAC 550
 CTGTACCTGC TGCTGGCGGT CTTAAAGGAG TGGAAATCCT CACCAGGCTC 600
 CAACCTAGGG TCTGCACCCA TGGCACTGGA GCGGAACCTG TCAGAGCTGC 650
 10 AGCTTCAGGC AGCTGTGGAA GCGACAGGGG ACCTGGAGGT AGACTGCTAC 700
 CGGGCACCTT GCTGTGAGAG CCAGGAAGAA CTGGCCCTGC AGCACTTGGT 750
 GAAGGAGAAG CTGCTGGCTC AGCTGGAGGA GAAGCTGCGT GTGTTTGCAA 800
 ACATTGTTGC TGTCTCAAC AAGGAAGTGG AGGCTTCCCA CCTGGCACTG 850
 GCCGCCTCCA TCCACCAGAG CCAGTTGGAC CGAGAGCACC TCCTGAGCTT 900
 15 GGAGCAGAGG GTGGTGGAAT TACAGCAAAC CCTGGCTCAA AAAGACCAGG 950
 TCCTGGGCAA GCTTGAGCAC AGTCTGCGAC TCATGGAGGA GGCATCCTTT 1000
 GATGGTACTT TCCTGTGGAA GATCACCAAT GTCACCAAGC GGTGCCACGA 1050
 GTCAGTGTGT GGCCGGACTG TCAGCCTCTT CTCTCCAGCT TTCTACACTG 1100
 CCAAGTATGG TTACAAGTTG TGCCTGCGCT TGTACCTGAA CGGGGATGGC 1150
 20 TCAGGCAAGA AGACCCACCT GTCCCTCTTC ATCGTGATCA TGAGAGGAGA 1200
 ATACGATGCT CTCCTGCCCT GGCCTTTCAG GAACAAGGTC ACCTTTATGC 1250
 TACTTGACCA GAACAACCGA GAGCATGCTA TTGATGCCTT CCGGCCTGAC 1300
 CTGAGCTCAG CTCCTTCCA GCGGCCACAG AGTGAGACCA ACGTGGCCAG 1350

CGGCTGCCCC CTCTTCTTCC CCCTCAGCAA GCTGCAGTCA CCCAAGCAGC 1400

CCTACGTCAA AGATGACACA ATGTTCTTCA AATGCATTGT GGACACTAGT 1450

GCTTAGGGAT GGGGGGAGGG GGTGTCTCCT GACAGAACCA GCTTAGACTG 1500

GGGGACTTAG CTAGACAGCC AGGCCCTGCC TGCCCTTGGA GCCCACAGCC 1550

5 CACGACAAGG AGGAGCCAAG GCTGGCATGA CTTTCAGCGCC ACAGCATGCT 1600

GGTTATGGCT GATGTGAGGC TGGAGAAACG TGTGCGTACA GAGACAGAGT 1650

GGAGGAGAAG ACAGAAGTGC TCTTTTCACA CAGACTACAC GACACCAGGA 1700

GGCCAGCATG CCAGCAGCTT CTGAATGTTG AGACCAGCCT AGATCAGGAT 1750

GAAAAGAGCC AGGCCTGAGG CTTGGACATT GAGCCAAGGC TATGGGGCCT 1800

10 AAGTGGAGGG GCACTCCTAC CAGGACATTC TCTCGAGGTC AGGGCATAAC 1850

TGGAAAAATG CCCCCATCTC TCTGTTTCTG CTCAAACTA GAACCACAGG 1900

GCAGAAGGGT CAGACATTAA TGTGAATTTA ACCTGCCCTG GACTGAGTTC 1950

CTATGTTAAC AGACACGCAA ACAGGTAAAC CCAGAACTG CCCTGGGAAA 2000

TGCTTTCTGG CTGCATCTGG AGATCTTTGA TGTTTTACC GACAAAACAA 2050

15 ATAACAAAAG CCTTGAATTG CAAAAAAAAA AAAAAAAAAA 2088

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 409 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Ala	Ser	Ser	Ser	Ala	Pro	Asp	Glu	Asn	Glu	Phe	Gln	Phe	Gly
1				5					10					15
Cys	Pro	Pro	Ala	Pro	Cys	Gln	Asp	Pro	Ser	Glu	Pro	Arg	Val	Leu
25				20					25					30
Cys	Cys	Thr	Ala	Cys	Leu	Ser	Glu	Asn	Leu	Arg	Asp	Asp	Glu	Asp
				35					40					45
Arg	Ile	Cys	Pro	Lys	Cys	Arg	Ala	Asp	Asn	Leu	His	Pro	Val	Ser
				50					55					60
Pro	Gly	Ser	Pro	Leu	Thr	Gln	Glu	Lys	Val	His	Ser	Asp	Val	Ala
30				65					70					75

	Glu	Ala	Glu	Ile	Met	Cys	Pro	Phe	Ala	Gly	Val	Gly	Cys	Ser	Phe	80	85	90
	Lys	Gly	Ser	Pro	Gln	Ser	Met	Gln	Glu	His	Glu	Ala	Thr	Ser	Gln	95	100	105
5	Ser	Ser	His	Leu	Tyr	Leu	Leu	Leu	Ala	Val	Leu	Lys	Glu	Trp	Lys	110	115	120
	Ser	Ser	Pro	Gly	Ser	Asn	Leu	Gly	Ser	Ala	Pro	Met	Ala	Leu	Glu	125	130	135
10	Arg	Asn	Leu	Ser	Glu	Leu	Gln	Leu	Gln	Ala	Ala	Val	Glu	Ala	Thr	140	145	150
	Gly	Asp	Leu	Glu	Val	Asp	Cys	Tyr	Arg	Ala	Pro	Cys	Cys	Glu	Ser	155	160	165
	Gln	Glu	Glu	Leu	Ala	Leu	Gln	His	Leu	Val	Lys	Glu	Lys	Leu	Leu	170	175	180
15	Ala	Gln	Leu	Glu	Glu	Lys	Leu	Arg	Val	Phe	Ala	Asn	Ile	Val	Ala	185	190	195
	Val	Leu	Asn	Lys	Glu	Val	Glu	Ala	Ser	His	Leu	Ala	Leu	Ala	Ala	200	205	210
20	Ser	Ile	His	Gln	Ser	Gln	Leu	Asp	Arg	Glu	His	Leu	Leu	Ser	Leu	215	220	225
	Glu	Gln	Arg	Val	Val	Glu	Leu	Gln	Gln	Thr	Leu	Ala	Gln	Lys	Asp	230	235	240
	Gln	Val	Leu	Gly	Lys	Leu	Glu	His	Ser	Leu	Arg	Leu	Met	Glu	Glu	245	250	255
25	Ala	Ser	Phe	Asp	Gly	Thr	Phe	Leu	Trp	Lys	Ile	Thr	Asn	Val	Thr	260	265	270
	Lys	Arg	Cys	His	Glu	Ser	Val	Cys	Gly	Arg	Thr	Val	Ser	Leu	Phe	275	280	285
30	Ser	Pro	Ala	Phe	Tyr	Thr	Ala	Lys	Tyr	Gly	Tyr	Lys	Leu	Cys	Leu	290	295	300
	Arg	Leu	Tyr	Leu	Asn	Gly	Asp	Gly	Ser	Gly	Lys	Lys	Thr	His	Leu	305	310	315
	Ser	Leu	Phe	Ile	Val	Ile	Met	Arg	Gly	Glu	Tyr	Asp	Ala	Leu	Leu	320	325	330
35	Pro	Trp	Pro	Phe	Arg	Asn	Lys	Val	Thr	Phe	Met	Leu	Leu	Asp	Gln	335	340	345
	Asn	Asn	Arg	Glu	His	Ala	Ile	Asp	Ala	Phe	Arg	Pro	Asp	Leu	Ser	350	355	360
40	Ser	Ala	Ser	Phe	Gln	Arg	Pro	Gln	Ser	Glu	Thr	Asn	Val	Ala	Ser	365	370	375
	Gly	Cys	Pro	Leu	Phe	Phe	Pro	Leu	Ser	Lys	Leu	Gln	Ser	Pro	Lys	380	385	390
	His	Ala	Tyr	Val	Lys	Asp	Asp	Thr	Met	Phe	Leu	Lys	Cys	Ile	Val	395	400	405
45	Asp	Thr	Ser	Ala												409		

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2121 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GCGCGAAGAC CGTTGGGGCT TTGTGGTGTG TGGGGGTTGT AACTCACATG 50
GCTGCAGCCA GTGTGACTTC CCCTGGCTCC CTAGAACTGC TACAGCCTGG 100
10 CTTCTCCAAG ACCCTCCTGG GGACCAGGTT AGAAGCCAAG TACCTCTGTT 150
CAGCCTGCAA AAACATCCTG CGGAGGCCTT TCCAGGCCCA GTGTGGGCAC 200
CGCTACTGCT CTTTCTGCCT GACCAGCATC CTCAGCTCTG GGCCCCAGAA 250
CTGTGCTGCC TGTGTCTATG AAGGCCTGTA TGAAGAAGGC ATTTCTATTT 300
TAGAGAGTAG TTCGGCCTTT CCAGATAACG CTGCCCCGAG AGAGGTGGAG 350
15 AGCCTGCCAG CTGTCTGTCC CAATGATGGA TGCACTTGGA AGGGGACCTT 400
GAAAGAATAC GAGAGCTGCC ACGAAGGACT TTGCCCATTC CTGCTGACGG 450
AGTGTCTGCT ATGTAAAGGC CTGGTCCGCC TCAGCGAGAA GGAGCACCAC 500
ACTGAGCAGG AATGCCCCAA AAGGAGCCTG AGCTGCCAGC ACTGCAGAGC 550
ACCCTGTAGC CACGTGGACC TGGAGGTACA CTATGAGGTC TGCCCCAAGT 600
20 TTCCCTTAAC CTGTGATGGC TGTGGCAAGA AGAAGATCCC TCGGGAGACG 650
TTTCAGGACC ATGTTAGAGC ATGCAGCAAA TGCCGGGTTT TCTGCAGATT 700
CCACACCGTT GGCTGTTTCA AGATGGTGGG GACTGAGAAC CTGCAGGATC 750
ATGAGCTGCA GCGGCTACGG GAACACCTAG CCCTACTGCT GAGCTCATTC 800
TTGGAGGCCC AAGCCTCTCC AGGAACCTTG AACCAGGTGG GGCCAGAGCT 850
25 ACTCCAGCGG TGCCAGATTT TGGAGCAGAA GATAGCAACC TTTGAGAACA 900
TTGTCTGCGT CTTGAACCGT GAAGTAGAGA GGGTAGCAGT GACTGCAGAG 950
GCTTGTAGCC GGCAGCACCG GCTAGACCAG GACAAGATTG AGGCCCTGAG 1000

TAACAAGGTG CAACAGCTGG AGAGGAGCAT CGGCCTCAAG GACCTGGCCA 1050
 TGGCTGACCT GGAGCAGAAG GTCTCCGAGT TGGAAGTATC CACCTATGAT 1100
 GGGGTCTTCA TCTGGAAGAT CTCTGACTTC ACCAGAAAGC GTCAGGAAGC 1150
 CGTAGCTGGC CGGACACCAG CTATCTTCTC CCCAGCCTTC TACACAAGCA 1200
 5 GATATGGCTA CAAGATGTGT CTACGAGTCT ACTTGAATGG CGACGGCACT 1250
 GGGCGGGGAA CTCATCTGTC TCTCTTCTTC GTGGTGATGA AAGGCCCCAA 1300
 TGATGCTCTG TTGCAGTGGC CTTTAAATCA GAAGGTAACA TTGATGTTGC 1350
 TGGACCATAA CAACCGGGAG CATGTGATCG ACGCATTGAG GCCCGATGTA 1400
 ACCTCGTCCT CCTTCCAGAG GCCTGTCACT GACATGAACA TCGCCAGTGG 1450
 10 CTGCCCCCTC TTCTGCCCTG TGTCCAAGAT GGAGGCCAAG AATTCCTATG 1500
 TGCGGGATGA TGCGATCTTC ATCAAAGCTA TTGTGGACCT AACAGGACTC 1550
 TAGCCACCCC TGCTAAGAAT AGCAGCTCAG TGAGGAGCTG TCACATTAGG 1600
 CCAGCCAGGC CCTGCCACAC ACGGGTGGGC AGGCTTGGTG TAAATGCTGG 1650
 GGAGGGCCTC AGCCTAGAGC CAATCACCAT CACACAGAAA GGCAGGAAGA 1700
 15 AGCCTCCAGT TGGCCTTCAG CTGGCAAACCT GAGTTGGACG GTCCACTGAG 1750
 CTCAAGGGCC TGGTGGAGCC CGCTGGGGAG CTTCTCAGCT TTCCAATAGG 1800
 AAAGCTCCTG CTGTCTCCTC TGTCTGGGGA AGGGAGAGAC CTGTAGGTGG 1850
 GTGCTCAGAA AGGGCCTCTC CAGAGAGAGT CTCAAGAGCT GCAGCAGGAG 1900
 CAAAGTGACT GGCCTTCCCC ACCCCATCCT TTGGAAAAGA GGTAGCGGCT 1950
 20 ACACAGGAGA AGGCATGCGC CTGCAGGGTG TAGCCCAAGA GAGAAGCTCT 2000
 CTGAGACATA GGCCCTCACT GGAGAAGGGC CTGCCTGGGC TGCACAGCCT 2050
 TGCCAGGTGG CCTGTATGGG GGAGAAGTGA TTAAATGTTG AGATGTCACA 2100
 CGACAAAAAA AAAAAAAAAA A 2121

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 501 amino acids

(B) TYPE: amino acid

5 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

	Met	Ala	Ala	Ala	Ser	Val	Thr	Ser	Pro	Gly	Ser	Leu	Glu	Leu	Leu	1	5	10	15
10	Gln	Pro	Gly	Phe	Ser	Lys	Thr	Leu	Leu	Gly	Thr	Arg	Leu	Glu	Ala	20	25	30	
	Lys	Tyr	Leu	Cys	Ser	Ala	Cys	Lys	Asn	Ile	Leu	Arg	Arg	Pro	Phe	35	40	45	
	Gln	Ala	Gln	Cys	Gly	His	Arg	Tyr	Cys	Ser	Phe	Cys	Leu	Thr	Ser	50	55	60	
15	Ile	Leu	Ser	Ser	Gly	Pro	Gln	Asn	Cys	Ala	Ala	Cys	Val	Tyr	Glu	65	70	75	
	Gly	Leu	Tyr	Glu	Glu	Gly	Ile	Ser	Ile	Leu	Glu	Ser	Ser	Ser	Ala	80	85	90	
20	Phe	Pro	Asp	Asn	Ala	Ala	Arg	Arg	Glu	Val	Glu	Ser	Leu	Pro	Ala	95	100	105	
	Val	Cys	Pro	Asn	Asp	Gly	Cys	Thr	Trp	Lys	Gly	Thr	Leu	Lys	Glu	110	115	120	
	Tyr	Glu	Ser	Cys	His	Glu	Gly	Leu	Cys	Pro	Phe	Leu	Leu	Thr	Glu	125	130	135	
25	Cys	Pro	Ala	Cys	Lys	Gly	Leu	Val	Arg	Leu	Ser	Glu	Lys	Glu	His	140	145	150	
	His	Thr	Glu	Gln	Glu	Cys	Pro	Lys	Arg	Ser	Leu	Ser	Cys	Gln	His	155	160	165	
30	Cys	Arg	Ala	Pro	Cys	Ser	His	Val	Asp	Leu	Glu	Val	His	Tyr	Glu	170	175	180	
	Val	Cys	Pro	Lys	Phe	Pro	Leu	Thr	Cys	Asp	Gly	Cys	Gly	Lys	Lys	185	190	195	
	Lys	Ile	Pro	Arg	Glu	Thr	Phe	Gln	Asp	His	Val	Arg	Ala	Cys	Ser	200	205	210	
35	Lys	Cys	Arg	Val	Leu	Cys	Arg	Phe	His	Thr	Val	Gly	Cys	Ser	Glu	215	220	225	
	Met	Val	Glu	Thr	Glu	Asn	Leu	Gln	Asp	His	Glu	Leu	Gln	Arg	Leu	230	235	240	
40	Arg	Glu	His	Leu	Ala	Leu	Leu	Leu	Ser	Ser	Phe	Leu	Glu	Ala	Gln	245	250	255	
	Ala	Ser	Pro	Gly	Thr	Leu	Asn	Gln	Val	Gly	Pro	Glu	Leu	Leu	Gln	260	265	270	
	Arg	Cys	Gln	Ile	Leu	Glu	Gln	Lys	Ile	Ala	Thr	Phe	Glu	Asn	Ile	275	280	285	
45	Val	Cys	Val	Leu	Asn	Arg	Glu	Val	Glu	Arg	Val	Ala	Val	Thr	Ala	290	295	300	

Glu Ala Cys Ser Arg Gln His Arg Leu Asp Gln Asp Lys Ile Glu
 305 310 315
 Ala Leu Ser Asn Lys Val Gln Gln Leu Glu Arg Ser Ile Gly Leu
 320 325 330
 5 Lys Asp Leu Ala Met Ala Asp Leu Glu Gln Lys Val Ser Glu Leu
 335 340 345
 Glu Val Ser Thr Tyr Asp Gly Val Phe Ile Trp Lys Ile Ser Asp
 350 355 360
 10 Phe Thr Arg Lys Arg Gln Glu Ala Val Ala Gly Arg Thr Pro Ala
 365 370 375
 Ile Phe Ser Pro Ala Phe Tyr Thr Ser Arg Tyr Gly Tyr Lys Met
 380 385 390
 Cys Leu Arg Val Tyr Leu Asn Gly Asp Gly Thr Gly Arg Gly Thr
 395 400 405
 15 His Leu Ser Leu Phe Phe Val Val Met Lys Gly Pro Asn Asp Ala
 410 415 420
 Leu Leu Gln Trp Pro Phe Asn Gln Lys Val Thr Leu Met Leu Leu
 425 430 435
 20 Asp His Asn Asn Arg Glu His Val Ile Asp Ala Phe Arg Pro Asp
 440 445 450
 Val Thr Ser Ser Ser Phe Gln Arg Pro Val Ser Asp Met Asn Ile
 455 460 465
 Ala Ser Gly Cys Pro Leu Phe Cys Pro Val Ser Lys Met Glu Ala
 470 475 480
 25 Lys Asn Ser Tyr Val Arg Asp Asp Ala Ile Phe Ile Lys Ala Ile
 485 490 495
 Val Asp Leu Thr Gly Leu
 500 501

(2) INFORMATION FOR SEQ ID NO:5:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 44 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

35 Asp Leu Leu Cys Pro Ile Cys Met Gln Ile Ile Lys Asp Ala Phe
 1 5 10 15
 Leu Thr Ala Cys Gly His Ser Phe Cys Tyr Met Cys Ile Ile Thr
 20 25 30
 40 His Leu Arg Asn Lys Ser Asp Cys Pro Cys Cys Ser Gln His
 35 40 44

(2) INFORMATION FOR SEQ ID NO:6:

- 45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 47 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Glu Leu Ser Cys Ser Ile Cys Leu Glu Pro Phe Lys Glu Pro Val
 1 5 10 15
 Thr Thr Pro Cys Gly His Asn Phe Cys Gly Ser Cys Leu Asn Glu
 20 25 30
 5 Thr Trp Ala Val Gln Gly Ser Pro Tyr Leu Cys Pro Gln Cys Arg
 35 40 45
 Ala Val
 47

(2) INFORMATION FOR SEQ ID NO:7:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 44 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

15 Leu Leu Arg Cys His Ile Cys Lys Asp Phe Leu Lys Val Pro Val
 1 5 10 15
 Leu Thr Pro Cys Gly His Thr Phe Cys Ser Leu Cys Ile Arg Thr
 20 25 30
 20 His Leu Asn Asn Gln Pro Asn Cys Pro Leu Cys Leu Phe Glu
 35 40 44

(2) INFORMATION FOR SEQ ID NO:8:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 44 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Ala Phe Arg Cys His Val Cys Lys Asp Phe Tyr Asp Ser Pro Met
 1 5 10 15
 30 Leu Thr Ser Cys Asn His Thr Phe Cys Ser Leu Cys Ile Arg Arg
 20 25 30
 Cys Leu Ser Val Asp Ser Lys Cys Pro Leu Cys Arg Ala Thr
 35 40 44

(2) INFORMATION FOR SEQ ID NO:9:

- 35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 45 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

40 Ser Ile Ser Cys Gln Ile Cys Glu His Ile Leu Ala Asp Pro Val
 1 5 10 15
 Glu Thr Asn Cys Lys His Val Phe Cys Arg Val Cys Ile Leu Arg
 20 25 30
 Cys Leu Lys Val Met Gly Ser Tyr Cys Pro Ser Cys Arg Tyr Pro
 35 40 45

45 (2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 45 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Glu Val Thr Cys Pro Ile Cys Leu Asp Pro Phe Val Glu Pro Val
 1 5 10 15
 Ser Ile Glu Cys Gly His Ser Phe Cys Gln Glu Cys Ile Ser Gln
 20 25 30
 10 Val Gly Lys Gly Gly Gly Ser Val Cys Ala Val Cys Arg Gln Arg
 35 40 45

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 46 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Glu Leu Met Cys Pro Ile Cys Leu Asp Met Leu Lys Asn Thr Met
 1 5 10 15
 20 Thr Thr Lys Glu Cys Leu His Arg Phe Cys Ser Asp Cys Ile Val
 20 25 30
 Thr Ala Leu Arg Ser Gly Asn Lys Glu Cys Pro Thr Cys Arg Lys
 35 40 45
 Lys
 25 46

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 50 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Glu Val Thr Cys Pro Ile Cys Leu Glu Leu Leu Lys Glu Pro Val
 1 5 10 15
 35 Ser Ala Asp Cys Asn His Ser Phe Cys Arg Ala Cys Ile Thr Leu
 20 25 30
 Asn Tyr Glu Ser Asn Arg Asn Thr Asp Gly Lys Gly Asn Cys Pro
 35 40 45
 Val Cys Arg Val Pro
 50

40 (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 47 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Glu Thr Thr Cys Pro Val Cys Leu Gln Tyr Phe Ala Glu Pro Met

1	5	10	15
Met Leu Asp Cys Gly His Asn Ile Cys Cys Ala Cys Leu Ala Arg	20	25	30
Cys Trp Gly Thr Ala Glu Thr Asn Val Ser Cys Pro Gln Cys Arg	35	40	45
Glu Thr	47		

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 48 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```

      Phe Gln Leu Cys Lys Ile Cys Ala Glu Asn Asp Lys Asp Val Lys
15       1              5             10            15
      Ile Glu Pro Cys Gly His Leu Met Cys Thr Ser Cys Leu Thr Ser
               20           25          30
      Trp Gln Glu Ser   Glu Gly Gln Gly Ser Ser Gly Cys Pro Phe Cys
                35         40        45
20    Arg Cys Glu
        48
```

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Gly Gly Phe Lys Leu Val Thr Cys Asp Phe Cys Lys Arg Asp Asp
 1 5 10 15

30 Ile Lys Lys Lys Glu Leu Glu Thr His Tyr Lys Thr Cys
 20 25 28

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 26 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

	Gln	Asp	Leu	Ala	Val	Cys	Asp	Val	Cys	Asn	Arg	Lys	Phe	Arg	His
	1				5					10					15
40	Lys	Asp	Tyr	Leu	Arg	Asp	His	Gln	Lys	Thr	His				
					20					25	26				

(2) INFORMATION FOR ~SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Thr Gly Lys Tyr Pro Phe Ile Cys Ser Glu Cys Gly Lys Ser Phe
1 5 10 15

5 Met Asp Lys Arg Tyr Leu Lys Ile His Ser Asn Val His
20 25 28

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids
(B) TYPE: amino acid
10 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Thr Gly Glu Lys Pro Tyr Thr Cys Thr Val Cys Gly Lys Lys Phe
1 5 10 15

15 Ile Asp Arg Ser Ser Val Val Lys His Ser Arg Thr His
20 25 28

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids
(B) TYPE: amino acid
20 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Arg Lys Lys Phe Pro His Ile Cys Gly Glu Cys Gly Lys Gly Phe
1 5 10 15

25 Arg His Pro Ser Ala Leu Lys Lys His Ile Arg Val His
20 25 28

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids
(B) TYPE: amino acid
30 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Ser Glu Glu Lys Pro Phe Glu Cys Glu Glu Cys Gly Lys Lys Phe
1 5 10 15

35 Arg Thr Ala Arg His Leu Val Lys His Gln Arg Ile His
20 25 28

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids
(B) TYPE: amino acid
40 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Pro Asn Glu Gln Met Ala Gln Cys Pro Ile Cys Gln Gln Phe Tyr
1 5 10 15

45 Pro Leu Lys Ala Leu Glu Lys Thr His Leu Asp Glu Cys
20 25 28

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Pro Asp Asp Gly Leu Val Ala Cys Pro Ile Cys Leu Thr Arg Met
1 5 10 15
Lys Glu Gln Gln Val Asp Arg His Leu Asp Thr Ser Cys
20 25 28

10 (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CCTTGTGCCT GCAGAGAGAA G 21

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
20 (A) LENGTH: 35 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

25 CTAGGTTAAC TTTCGGTGCT CCCCAGCAGG GTCTC 35

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
30 (A) LENGTH: 35 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CTAGGTTAAC TGGAGAAGGG GACCTGCTCG TCCTT 35

(2) INFORMATION FOR SEQ ID NO:26:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

CTAGGTTAAC TGCTGGCTTG GGAGGAGCAC TGTGA 35

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 35 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

CTAGGTTAAC TGCTCCCGGT GCTGGCCCGG GCCTC 35

10 (2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 34 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

CTAGGTTAAC TGCACTGGCC GAGCTCTCCA GGGA 34

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 15 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

25 GTGATGAGAA TTCAT 15

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 21 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

CGATGAATTC TCATCACTGC A 21

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 33 bases
(B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GATCGGATCC AAAAAGAAGC CTTGTGCCT GCA 33

5 (2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GCCTGGTTAA CTGGGC 16

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

20 GCNCCNATGG CNYTNGARC 19

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GCNCCNATGG CNYTNGARA 19

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

GCNCCNATGG CNYTNGARG 19

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 19 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

GYTCNARNGC CATNGGNGC 19

(2) INFORMATION FOR SEQ ID NO:37:

10 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

TYTCNARNGC CATNGGNGC 19

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 19 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

CYTCNARNGC CATNGGNGC 19

25 (2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 17 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

AARCAYGCNT AYG TNAA 17

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 17 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

TTNACRTANG CRTGYTT 17

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Ala Pro Met Ala Leu Glu Arg
1 5 7

10 (2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

Lys His Ala Tyr Val Lys
1 5 6

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

25 Pro Gly Ser Asn Leu Gly Ser
1 5 7

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

Lys Asp Asp Thr Met Phe Leu Lys
1 5 8

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 37 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

TCGATCGTCG ACCAAAAAGA AGCCCTCCTG CCTACAA 37

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:
- CTAGAGATCT CAGGGGTCAG GCCACTTT 28
- (2) INFORMATION FOR SEQ ID NO:47:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:
- CTAGAGATCT GTTAACTTTC GGTGCTCCCC AGCAGGGTCT C 41
- (2) INFORMATION FOR SEQ ID NO:48:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:
- CTAGAGATCT GTTAACTGGA GAAGGGGACC TGCTCGTCCT T 41
- (2) INFORMATION FOR SEQ ID NO:49:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:
- CTAGAGATCT GTTAACTGCT GGCTTGGGAG GAGCACTGTG A 41
- (2) INFORMATION FOR SEQ ID NO:50:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

TCGATCGTCG ACCAAAAAGA AGCCCTCCTG CCTACAA 37

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 28 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CTAGAGATCT CAGGGGTCAG GCCACTTT 28

10 (2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 34 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

TCGATCGTCG ACCGCCTCCA GCTCAGCCCC TGAT 34

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 31 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

25 GATCGGATCC GGAGACACAG ATTCCAGCCC C 31

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 35 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GATCGAATTC TTAACCTTTC GGTGCTCCCC AGCAG 35

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 30 bases
(B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

GATCGGATCC TTGTGGTGTG TGGGGGTTGT 30

5 (2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

CCTGGCTGGC CTAATGT 17

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

20 GATCGACTCG AGATGCCCAA GAAGAAGCGG AAGGTGGCTG CAGCCAGTGT 50

GACTTCCCCT 60

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

CTCTGGCGAA GAAGTCC 17

30 (2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

GATCGGATCC GCCTCCAGCT CAGCCCCTGA T 31

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 31 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

GATCGGATCC AGCCAGCAGC TTCTCCTTCA C 31

10 (2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 30 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

GATCGGATCC TTGTGGTGTG TGGGGGTTGT 30

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 30 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

25 GATCGGATCC GCTCAGGCTC TTTTGGGGCA 30

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 34 bases
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

GATCGAGTCG ACCAGTAGTT CGGCCTTTCA AGAT 34

(2) INFORMATION FOR SEQ ID NO:64:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 17 bases
(B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

CTCTGGCGAA GAAGTCC 17

5 (2) INFORMATION FOR SEQ ID NO:65:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

GATCGAGTCG ACCGTGGGGC CAGAGCTACT CCAG 34

(2) INFORMATION FOR SEQ ID NO:66:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

20 GTACGTGCGG CCGCCTACCA CCTGGTTCAA GGTTCTCTG 38

CLAIMS:


1. An isolated tumor necrosis receptor associated factor (TRAF) capable of specific association with the intracellular domain of a native type 2 TNF receptor (TNF-R2).
2. The TRAF of claim 1 that is murine.
- 5 3. The TRAF of claim 1 that is capable of specific association with the intracellular domain of a native human TNF-R2.
4. The TRAF of claim 1 that is capable of specific binding to the intracellular domain of a native human TNF-R2.
5. The TRAF of claim 1 that is native.
- 10 6. The TRAF of claim 5 in homodimeric form.
7. The TRAF of claim 5 associated with another TRAF to form a heterodimer.
8. The TRAF of claim 5 that is TRAF1 (SEQ. ID. NO: 2) or TRAF2 (SEQ. ID. NO: 4).
9. The TRAF of claim 1 which comprises a domain having at least about 50% sequence identity with the aa272-501 amino acid region of the TRAF2 amino acid sequence (SEQ. ID. NO: 4).
- 15 10. The TRAF of claim 1 encoded by nucleic acid molecule capable of hybridizing, under stringent conditions, to the complement of the nucleotide sequence encoding amino acids 272-501 of the TRAF2 amino acid sequence (SEQ. ID. NO: 4).
11. An isolated nucleic acid molecule comprising a nucleotide sequence encoding a TRAF of claim 1.
- 20 12. A vector comprising the nucleic acid molecule of claim 11 operably linked to control sequences recognized by a host cell transformed with the vector.
13. A host cell transformed with a vector of claim 12.
14. A molecule capable of disrupting the interaction of a TRAF and a native TNF-R2.
15. An antibody capable of specific binding to a native TRAF polypeptide.
- 25 16. A hybridoma cell line producing an antibody of claim 15.
17. A method of using a nucleic acid molecule encoding a TRAF comprising expressing such nucleic acid molecule in a cultured host cell transformed with a vector comprising such nucleic acid molecule operably linked to control sequences recognized by said host cell, and recovering the polypeptide encoded by said nucleic acid molecule from the host cell.
- 30 18. A method for producing a TRAF polypeptide comprising inserting into the DNA of a cell containing nucleic acid encoding said polypeptide a transcription modulatory element in sufficient proximity and orientation to the nucleic acid molecule to influence the transcription thereof.
19. A method of determining the presence of a TRAF polypeptide comprising hybridizing DNA encoding such polypeptide to a test sample nucleic acid and determining the presence of TRAF polypeptide DNA.
- 35 20. An isolated nucleic acid molecule encoding a fusion of an intracellular domain sequence of a native TNF-R2 to the DNA-binding domain of a transcriptional activator.
21. The nucleic acid molecule of claim 20, wherein said transcriptional activator is yeast GAL4.

22. An isolated nucleic acid molecule encoding a fusion of a TRAF to the activation domain of a transcriptional activator.
23. The nucleic acid molecule of claim 22, wherein said transcriptional activator is yeast GAL4.
24. A vector comprising the nucleic acid molecule of claim 20.
- 5 25. A vector comprising the nucleic acid molecule of claim 22.
26. An assay for identifying a factor capable of specific binding to the intracellular domain of a native TNF-R2, comprising
- (a) expressing nucleic acid molecules encoding a polypeptide comprising a fusion of an intracellular domain sequence of a native TNF-R2 to the DNA-binding domain of a transcriptional activator, and a fusion
- 10 of a candidate polypeptide factor to the activation domain of a transcriptional activator, in a single host cell carrying a reporter gene;
- (b) monitoring the binding of said candidate factor to the intracellular domain of TNF-R2 by detecting a signal of the molecule encoded by said reporter gene.
27. An assay for identifying a factor capable of specific association with the intracellular domain
- 15 of a native TNF-R2, comprising
- (a) expressing nucleic acid molecules encoding a polypeptide comprising a fusion of an intracellular domain sequence of a native TNF-R2 to the DNA-binding domain of a transcriptional activator, and a fusion of a candidate factor to the activation domain of a transcriptional activator, in a single host cell transfected with nucleic acid encoding a polypeptide factor capable of specific binding to said TNF-R2, and with nucleic
- 20 acid encoding a reporter gene; and
- (b) monitoring the association of said candidate factor with said TNF-R2 or with said polypeptide factor capable of specific binding to said TNF-R2 by detecting the signal of the polypeptide encoded by said reporter gene.
28. A method of amplifying a nucleic acid test sample, comprising priming a nucleic acid
- 25 polymerase reaction with nucleic acid encoding a TRAF polypeptide capable of specific association with the intracellular domain of a native TNF-R2.
29. A method for detecting a nucleic acid sequence coding for a polypeptide molecule which comprises all or part of a TRAF polypeptide or a related nucleic acid sequence, comprising contacting the nucleic acid sequence with a detectable marker which binds specifically to at least part of said nucleic acid
- 30 sequence, and detecting the marker so bound.
30. A method for the prevention or treatment of a pathological condition associated with a TNF biological activity mediated, fully or partially, by a TNF-R2, comprising administering to a patient in need a preventatively or therapeutically effective amount of a TRAF or a molecule capable of disrupting the interaction of a TRAF and said TNF-R2.
- 35 31. An assay for identifying a molecule capable of disrupting the association of a TRAF with the intracellular domain of a native TNF-R2, comprising contacting a cell coexpressing a native TNF-R2, a native TRAF polypeptide and a reporter gene with a candidate molecule, and monitoring the ability of said

candidate molecule to disrupt the association of said TRAF and TNF-R2 intracellular domain sequence by detecting the molecule encoded by the reporter gene.

32. As assay for identifying a molecule capable of disrupting the association of a TRAF with the intracellular domain of a native TNF-R2, comprising contacting a cell expressing 1. a fusion of an intracellular domain sequence of a native TNF-R2 to the DNA-binding domain of a transcriptional activator, 2. a fusion of a native TRAF polypeptide to the activation domain of said transcriptional activator, and 3. a reporter gene, with a candidate molecule, and monitoring the ability of said candidate molecule to disrupt the association of said TRAF and TNF-R2 intracellular domain sequence by detecting the molecule encoded by the reporter gene.
33. An assay for identifying an inhibitor of the interaction of a TRAF protein with CD40 comprising contacting recombinant host cells coexpressing a TRAF protein capable of direct or indirect binding of CD40, CD40 and a reporter gene the expression of which is dependent on the CD40:TRAF interaction, with candidate inhibitors and selecting a molecule which inhibits the expression of said reporter gene.
34. The assay of claim 33 wherein said TRAF protein is TRAF2.
35. The assay of claim 33 wherein said recombinant host cells coexpress TRAF1 and TRAF2.
36. The assay of claim 33 wherein said reporter gene is NF- κ B dependent.
37. The assay of claim 36 wherein said NF- κ B dependent reporter gene is an E-selectin-luciferase reporter gene construct.
38. The assay of claim 33 which is performed in the two-hybrid format.
39. An assay for identifying an inhibitor of the interaction of a TRAF protein with LMP1 comprising contacting recombinant host cells coexpressing a TRAF protein capable of direct or indirect binding of LMP1, LMP1 and a reporter gene the expression of which is dependent on the LMP1:TRAF interaction, with candidate inhibitors and selecting a molecule which inhibits the expression of said reporter gene.
40. The assay of claim 39 wherein said TRAF protein is TRAF2.
41. The assay of claim 39 wherein said recombinant host cells coexpress TRAF1 and TRAF2.
42. The assay of claim 39 wherein said reporter gene is NF- κ B dependent.
43. The assay of claim 42 wherein said NF- κ B dependent reporter gene is an E-selectin-luciferase reporter gene construct.
44. The assay of claim 39 which is performed in the two-hybrid format.
45. TRAF2(87-501) or a functional derivative thereof capable of inhibiting a biological activity mediated by TNF-R2, CD40 or LMP1.
46. The functional derivative of claim 45 which is capable of inhibiting TRN-R2-, CD4- or LMP1-mediated NF- κ B activation.

	Preimmune		Anti-mTNF-R2		Preimmune		Anti-mTNF-R2		Preimmune		Anti-mTNF-R2	
NF- κ B Probe	wt	wt	mt	mt	wt	wt	wt	wt	wt	wt	wt	wt
Competitor	-	-	-	-	mt	mt	AP-1					

**FIG. 1**




FIG. 2a


CT6
FIG. 2b

FIG. 3




FIG. 4






FIG. 5






FIG. 8




FIG. 9

```

1   CCCAGCCCGGTTCTCTGCCCCCAAGGACGCTACCGCCCAATGCGAGCAGAAGCGGCGCACAGATACAGAAAGT
74  GAGGCTCAGACATATTGAAGACCGTGTGACATAGGGTAGCCAAATGACAGTGTGAGAAAGTGACATTTACTCAAG
149 GCCACCCAGATATCCTGGAGGACCCAGAACCCCTGGAGATTCCCATCAGAAAGACCTTCTGCTCACCTTGAAACCCC

1   MetaIaSerSerAlaProAspGluAsnGluPheGlnPheGlyCysProProAlaProCysGlnAspPro
224 AAGATGGCCTCCAGCTCAGCCCTGATGAAACGAGTTTCAATTGGTTGCCCCCTGCTCCCTGCCAGGACCCCA

25  SerGluProArgValLeuCysCysThrAlaCysLeuSerGluAsnLeuArgAspGluAspArgIleCysPro
299 TCGGAGCCCCAGAGTTCTCTGCTGCACAGCCTGTCTCTGAGAACCTGAGAGATGATGAGGATCGGATCTGTCTCT

50  LysCysArgAlaAspAsnLeuHisProValSerProGlySerProLeuThrGlnGluLysValHisSerAspVal
374 AAATGCAGAGCAGACAACCTCCATCCTGTGAGCCAGGAAGCCCTCTGACTCAGGAGAAGGTTCACTCTGATGTA

75  AlaGluAlaGluIleMetCysProPheAlaGlyValGlyCysSerPheLysGlySerProGlnSerMetGlnGlu
449 GCTGAGGCTGAATCATGTGCCCTTTTGCAGGTGTGGCTGTTCCTTCAAGGGGAGCCACAAATCCATGCCAGGAG

100 HisGluAlaThrSerGlnSerSerHisLeuTyrLeuLeuLeuAlaValLeuLysGluTrpLysSerSerProGly
524 CATGAGGCTACCTCCAGTCTCCACCTGACCTGTGCTGCTGGCGGTCTTAAAGGAGTGGAAATCCTCACCGGC

125 SerAsnLeuGlySerAlaProMetAlaLeuGluArgAsnLeuSerGluLeuGlnLeuAlaValGluAla
599 TCCAAACCTAGGGTCTGCACCCATGGCACTGGAGCGGAACCTGTTCAGAGCTGCAGCTTCAGGCAGCTGTGGAGCGG

150 ThrGlyAspLeuGluValAlaAspCysTyrArgAlaProCysCysGluSerGlnGluGluLeuAlaLeuGlnHisLeu
674 ACAGGGGACCTGGAGGTAGACTGCTACCGGGCACCTTGCTGTGAGAGCCAGGAAGAACTGGCCCTGCAGCCTTG

175 ValLysGluLysLeuAlaGlnLeuGluGluLysLeuArgValPheAlaAsnIleValAlaValLeuAsnLys
749 GTGAAGGAGAGCTGCTGGCTCAGCTGGAGGAGAAGCTGCGTGTGTTTGCAAAACATTGTTGCTGTCTCAACAAG

200 GluValGluAlaSerHisLeuAlaLeuAlaSerIleHisGlnSerGlnLeuAspArgGluHisLeuLeuSer
824 GAAGTGGAGGCTTCCACCTGGCACCTGGCCGCTCCATCCACAGAGCCAGTGGACCGAGAGCACCTCCTGAGC

225 LeuGluGlnArgValValGluLeuGlnGlnThrLeuAlaGlnLysAspGlnValLeuGlyLysLeuGluHisSer
899 TTGGAGCAGAGGGGTGGTGGAAATTACAGCAAACCTGGCTCAAAAGACCAGGTCTCTGGGCAAGCTTGAGCACAGT

```

6/19

SUBSTITUTE SHEET (RULE 26)

FIG. 10a

✓

FIG. 10b

1 1 GCGCGAAGACCGTTGGGGCTTTGTGGTGTGGGGTTGTAACCTCACAATGGCTGCAGCCAGTGACCTCCCCCT
 10 GlySerLeuGluLeuLeuGlnProGlyPheSerLysThrLeuLeuGlyThrArgLeuGluAlaLysTyrLeuCys
 75 GGCTCCCTAGAACTGCTACAGCCTGGCTTCTCCAAGACCCCTCTGGGACCAAGTTAGAAGCCCAAGTACCTCTGT
 35 SerAlaCysLysAsnIleLeuArgArgProPheGlnAlaGlnCysGlyHisArgTyrCysSerPheCysLeuThr
 150 TCAGCCTGCAAAACATCCTGCGGAGGCCCTTTCCAGGCCCAGTGTGGCACCCGCTACTGCTCCTTCTGCTGACC
 60 SerIleLeuSerSerGlyProGlnAsnCysAlaAlaCysValTyrGluGlyLeuTyrGluGlyIleSerIle
 225 AGCATCCTCAGCTCTGGGCCCCAGAACTGTGCTGCTGTGCTATGAAGGCCCTGTATGAAGAAGGCATTTCTATT
 85 LeuGluSerSerAlaPheProAspAsnAlaAlaArgArgGluValGluSerLeuProAlaValCysProAsn
 300 TTAGAGAGTAGTTCGGCCTTTCCAGATAACGCTGCCCGCAGAGAGGTGGAGAGCCTGCCAGCTGTCTGCCCAAT
 110 AspGlyCysThrTrpLysGlyThrLeuLysGluTyrGluSerCysHisGluGlyLeuCysProPheLeuLeuThr
 375 GATGGATGCACCTTGAAGGGGACCTTGAAAGAAATACGAGAGCTGCCACGAAGGACTTTGCCCATTTCTGCTGACG
 135 GluCysProAlaCysLysGlyLeuValArgLeuSerGluLysGluHisThrGluGlnGluCysProLysArg
 450 GAGTGTCCTGTCATGTAAAGGCCCTGGTCCGCTCAGCGAGAAGGAGCACCACTGAGCAGGAATGCCCCAAAGG
 160 SerLeuSerCysGlnHisCysArgAlaProCysSerHisValAspLeuGluValHisTyrGluValCysProLys
 525 AGCCTGAGCTGCCAGCACTGCAGAGCACCCCTGTAGCCACGTGGACCTGGAGGTACACTATGAGGTCTGCCCAAG
 185 PheProLeuThrCysAspGlyCysGlyLysLysIleProArgGluThrPheGlnAspHisValArgAlaCys
 600 TTTCCCTTAACCTGTGATGGCTGTGGCAAGAAGAAGATCCCTCGGGAGACGTTTCAGGACCATGTTAGAGCATGC
 210 SerLysCysArgValLeuCysArgPheHisThrValGlyCysSerGluMetValGluThrGluAsnLeuGlnAsp
 675 AGCAATGCGGGTCTCTGCAGATTCCACACCGTTGGCTGTTTCAGAGATGGTGAGACTGAGAACCCTGCAGGAT
 235 HisGluLeuGlnArgLeuArgGluHisLeuAlaLeuLeuSerSerPheLeuGluAlaGlnAlaSerProGly
 750 CATGAGCTGCAGCGGCTACGGGAACACCTAGCCCTACTGCTGAGCTCATTTCTTGGAGGCCCAAGCCTCTCCAGGA
 260 ThrLeuAsnGlnValGlyProGluLeuLeuGlnArgCysGlnIleLeuGluGlnLysIleAlaThrPheGluAsn
 825 ACCTTGAAACCGGTGGGGCCAGAGCTACTCCAGCGGTGCCAGATTTTGGAGCAGAGATAGCAACCTTTGAGAAC

8/19

SUBSTITUTE SHEET (RULE 26)

FIG. 11a

```

285 I leValCysValLeuAsnArgGluValGluArgValAlaValThrAlaGluAlaCysSerArgGlnHisArgLeu
900 ATTGTCCTGCGTCTTTGAACCGTGAAGTAGAGAGGGTAGCAGTACTGACAGAGGCTTGTAGCCGGCAGCACCGGCTA

310 AspGlnAspLysIleGluAlaLeuSerAsnLysValGlnGlnLeuGluArgSerIleGlyLeuLysAspLeuAla
975 GACCAGGACAAGATTGAGGCCCTGAGTAACAAGGTGCAACAGCTGGAGAGGAGCATCGGCCTCAAGGACCTGGCC

335 MetAlaAspLeuGluGlnLysValSerGluLeuGluValSerThrTyrAspGlyValPheIleTrpLysIleSer
1050 ATGGCTGACCTGGAGCAGAAGGCTCTCCGAGTTGGAAGTATCCACCTATGATGGGGTCTTCATCTGGGAAGATCTCT

360 AspPheThrArgLysArgGlnGluAlaValAlaGlyArgThrProAlaIlePheSerProAlaPheTyrThrSer
1125 GACTTCACCAGAAAGCGTCAGGAAGCCGTAGCTGGCCGGACACCAAGCTATCTTCTCCCCAGCCTTCTACACAAGC

385 ArgTyrGlyTyrLysMetCysLeuArgValTyrLeuAsnGlyAspGlyThrGlyArgGlyThrHisLeuSerLeu
1200 AGATATGGCTACAAGATGTGTCTACGAGTCTACTTGAATGGCGACGGCACTGGCGGGGAACCTCATCTGCTCTC

410 PhePheValValMetLysGlyProAsnAspAlaLeuLeuGlnTrpProPheAsnGlnLysValThrLeuMetLeu
1275 TTCTTCGTGGTGATGAAGGCCCAATGATGCTCTGTGTGAGTGGCCTTTTAAATCAGAAGGTAAACATTGATGTTG

435 LeuAspHisAsnAsnArgGluHisValIleAspAlaPheArgProAspValThrSerSerSerPheGlnArgPro
1350 CTGGACCATAACAACCGGGAGCATGTGATCGACGCAATTCAGGCCCGATGTAAACCTCGTCTCTCCAGAGGCCCT

460 ValSerAspMetAsnIleAlaSerGlyCysProLeuPheCysProValSerLysMetGluAlaLysAsnSerTyr
1425 GTCAGTGACATGAACATCGCCAGTGGCTGCCCCCTCTTCTGCCCTGTGTCCAAGATGGAGGCCAAGAAATTCCTAT

485 ValArgAspAspAlaIlePheIleLysAlaIleValAspLeuThrGlyLeu
1500 GTGCGGGATGATGCGATCTTCATCAAAGCTATTGTGGACCTAACAGGACTCTAGCCACCCCTGCTAAGAATAGCA
1575 GCTCAGTGAGGAGCTGTACATTAGGCCAGCCAGGCCCTGCCACACACGGGTGGGCAGGCTTGGTGTAAATGCTG
1650 GGGAGGGCCCTCAGCCTAGAGCCAATCACCATCACAGAAAGGACGGAAGCTCCAGTTGGCCTTCAGCTGG
1725 CAAACTGAGTTGGACGGTCCACTGAGCTCAAGGGCTGGTGGAGCCGCTGGGGAGCTTCTCAGCTTCCCAATAG
1800 GAAAGCTCCTGCTGCTCTGCTGGGAAGGGAGAGACTGTAGGTGGTGTCTAGAAAGGGCCTCTCCAGA
1875 GAGAGTCTCAAGAGCTGCAGCAGGAGCAAGTACTGGCTTCCCAACCCCTCTTGGAAAGAGGTAAGCGGC
1950 TACACAGGAGAAAGGCAATGCGCCTGCAGGTGTAGCCCAAGAGAGAAGCTCTCTGAGACATAGGCCCTCAGTGGAG
2025 AAGGGCCTGCCCTGGGCTGCACAGCCTTGCCAGGTGGCTGTATGGGGGAGAAGTGATTAATGTTGAGATGTCAC
2100 ACGACAAAAAATAAAAAA

```

FIG. 11b

FIG. 12a

TRAF2	(mouse)	31	KYLCSACKNILRRPQA QCGHRYCSFCLTSI	LSS	GPQNC AACVYE
COP1	(A. thaliana)	49	DLLCPICMQIIKDAFLT ACGHSFCYMCIIITH	LRN	KSDCPCCSQH
EFP	(human)	10	ELSCSICLEPFKEPVTT PCGHNF CGSCLNETWA VQG	SPYLCPQ	CRAV
RAD-18	(S. cerevisiae)	25	LLRCHICKDFLKVPLT PCGHTFCSLCIRTH	LNN	QPNCPCLLFE
UVS-2	(N. crassa)	31	APRCHVCKDFYDSPMLT SCNHTFCSLCIRRC	LSV	DSK CPLCRAT
RAG-1	(human)	290	SISCQICEHILADPVET NCKHVF CRVCILRC	LKV	MGSYCPSCRYP
SS-A/Ro	(human)	13	EVTCPICLDPFVEPVSIECGHSFCQECISQV	GKG	GGSVCAVCRQR
RING1	(human)	16	ELMCPICLDMLKNTWTTKECLHRFCSDCIVTA	LRS	GNKECPTCRKK
RFT-1	(mouse)	12	EVTCPICLELLKEPVA DCNH SFCRA C I T L N Y E S N R N T D G K G N C P V C R V P		
RFP	(human)	13	ETTCPVCLQYFAEPMML DCGHNICCA CLARCWGT A	ETNVSCPQ	CRET
c-cbl	(human)	378	FQLCKICAENDKDVKIE PCGHLMCTSC LTS WQESEGO	GSSGC	PPFCRCE

consensus

X11-12

X10-16

---C--C-----C-H--C--C-----C--C---
 X11-12 X10-16

FIG. 12b

TRAF2	(mouse)	157	CPKRSLS C QH C RAPCSHVDLEV H YE VC
		182	PKFPLT C DG C GKKKIPRET F QD H VR AC
DG17	(<i>D. discoideum</i>)	171	GGFKLVT C DF C KRDDIKKKELET H YK TC
TFIIIA	(<i>X. laevis</i>)	189	QD LAV C DV C NRKFRHKDYLRD H QK TH
XLCOF14	(<i>X. laevis</i>)	1	TGKYPFI C SE C GKSFMDKRYLKI H SN VH
XFIN	(<i>X. laevis</i>)	1225	TGEKPYT C TV C GKKFIDRSSVVK H SR TH
ZFY1/2	(mouse)	521	RKKFPHI C GE C GKGFRHPSALKK H IR VH
MFG2	(mouse)	293	SEEKPF E CE E C G KKFRTARHLVK H QR IH
RAD18	(<i>S. cerevisiae</i>)	183	PNEQMAQ C PI C QQFYPLKALEKT H LD EC
UVS-2	(<i>N. crassa</i>)	182	PDDGLVA C PI C LTRM KEQQVDR H LDTSC

TRAF2 1 MAAASVTSPGSLELLQPGFSKTLTGTRLEAKYLC SACKNILRRPFOAQCG
 TRAF2 51 HRYCSFCLTSILSSGPQNCAACVYEGLYEEGISILESSSAFPD NAARREV
 TRAF2 101 ESLPAVCPNDGCTWKGT LKEYESCHEGLCPFLLTECPACKGLVRLSEKEH
 TRAF1 1 MASSSAPDENEFOFGCPPA
 TRAF2 151 HTEQEC PK R S L S C Q H C R A P C S H V D L E V H Y E V C P K F P L T C D G C G K K K I P R E
 TRAF1 20 P C O D P S E P R V L C C T A C L S E N L R D D E D R I C P K C R A D N L H P V S P G S P L T Q E
 TRAF2 201 T F Q D H V R A C S K C R V L C R F H T V G C S E M V E T E N L Q D H E L O R L R E H L A L L L S S
 TRAF1 69 K V H S D V - - - A E A E I M C P F A G V G C S F K G S P Q S M Q E H E A T S Q S S H L Y L L L A V
 TRAF2 251 F L E A Q A S P G T L N Q V G P E L L Q R
 TRAF1 116 L K E W K S S P G S N L G S A P M A L E R N L S E L Q L Q A A V E A T G D L E V D C Y R A P C C E S
 TRAF2 272 C Q I L E Q K I A T F E N I V C V L N R E V E R V A V T A E A C S R Q H
 TRAF1 166 Q E E L A L O H L V K E K L L A Q L E E K L R V F A N I V A V L N K E V E A S H L A L A A S I H Q S
 TRAF2 308 R L D O D K I E A L S N K V Q Q L E R S I G L K D L A M A D L E O K V S E L E V S T Y D G V F I W K
 TRAF1 216 Q L D R E H L L S L E Q R V V E L Q O T L A Q K D O V L G K L E H S L R L M E E A S F D G T F L W K
 TRAF2 358 I S D F T R K R Q E A V A G R T P A I F S P A F Y T S R Y G Y K M C L R V Y L N G D G T G R G T H L
 TRAF1 266 I T N V T K R C H E S V C G R T V S L F S P A F Y T A K Y G Y K L C L R L Y L N G D G S G K K T H L
 TRAF2 408 S L F F V V M K G P N D A L L Q W P F N Q K V T L M L L D H N N R E H V I D A F R P D V T S S S F Q
 TRAF1 316 S L F I V I M R G E Y D A L L P W P F R N K V T F M L L D Q N N R E H A I D A F R P D L S S A S F Q
 TRAF2 458 R P V S D M N I A S G C P L F C P V S K M E - A K N S Y V R D D A I F I K A I V D L T G L
 TRAF1 366 R P Q S E T N V A S G C P L F F P L S K L Q S P K H A Y V K D D T M F L K C I V D T S A

FIG. 13




FIG. 14a




FIG. 14b




FIG. 15a




FIG. 15b




FIG. 16

FIG. 18a





FIG. 18b



Transfected DNA

18/19

SUBSTITUTE SHEET (RULE 26)

FIG. 19a





FIG. 19b



Transfected DNA

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/US 95/06639

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 6 C12N15/12 C12N15/62 C12N15/81 C12N5/12 C12N1/19
 C07K14/715 C07K16/18 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CELL, vol. 76, no. 6, 25 March 1994 CELL PRESS, CAMBRIDGE, MA, US;; pages 959-962, C.A. SMITH ET AL. 'The TNF receptor superfamily of cellular and viral proteins: Activation, costimulation, and death' cited in the application the whole document --- -/--	1-46

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

19 September 1995

Date of mailing of the international search report

17. 10. 95

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Hornig, H

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 95/06639

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CELL, vol. 74, no. 5, 10 September 1993 CELL PRESS,CAMBRIDGE,MA,US;; pages 845-853, L.A. TARTAGLIA ET AL. 'A novel domain within the 55kd TNF receptor signals cell death' cited in the application the whole document ---	1-46
A	KEYSTONE SYMPOSIUM ON HEMATOPOIESIS, BRECKENRIDGE, COLORADO, USA, JANUARY 4-11, 1994. JOURNAL OF CELLULAR BIOCHEMISTRY SUPPLEMENT 0 (18 PART A). 1994. 5. ISSN: 0733-1959, ABSTRACT A011, GOEDDEL D V ET AL 'TNF receptor signal transduction.' see abstract ---	1-46
A	PROC. NATL.ACAD SCI., vol. 88, no. 21, 1 November 1991 NATL. ACAD SCI.,WASHINGTON,DC,US;; pages 9578-9582, C.T. CHIEN ET AL. 'The two hybrid system: A method to identify and clone genes for proteins that interact with a protein of interest' the whole document ---	20-46
A	GENES & DEVELOPMENT, vol. 7, no. 4, April 1993 CSH LABORATORY PRESS, NEW YORK,US, pages 555-569, T. DURFEE ET AL. 'The retinoblastoma protein associates with the protein phosphatase type 1 catalytic subunit' the whole document ---	20-46
P,X	CELL (CAMBRIDGE, MASS.) (1994), 78(4), 681-92 CODEN: CELLB5;ISSN: 0092-8674, 26 August 1994 ROTHE, MIKE ET AL 'A novel family of putative signal transducers associated with the cytoplasmic domain of the 75 kDa tumor necrosis factor receptor' cited in the application the whole document ---	1-13, 15-29

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/06639

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	J. BIOL. CHEM. (1994), 269(48), 30069-72 CODEN: JBCHA3;ISSN: 0021-9258, 2 December 1994 HU, HONG MING ET AL 'A novel RING finger protein interacts with the cytoplasmic domain of CD40' cited by the applicant the whole document ---	1-46
T	FEBS LETT. (1995), 358(2), 113-18 CODEN: FEBLAL;ISSN: 0014-5793, 23 January 1995 SATO, TAKAAKI ET AL 'A novel member of the TRAF family of putative signal transducing proteins binds to the cytosolic domain of CD40' the whole document ---	1-46
T	CELL (CAMBRIDGE, MASS.) (1995), 80(3), 389-99 CODEN: CELLB5;ISSN: 0092-8674, 10 February 1995 MOSIALOS, GEOGRE ET AL 'The Epstein-Barr virus transforming protein LMP1 engages signaling proteins for the tumor necrosis factor receptor family' cited by the applicant the whole document ---	1-46
T	SCIENCE (WASHINGTON D C) 267 (5203). 1995. 1494-1498. ISSN: 0036-8075, 10 March 1995 CHENG G ET AL 'Involvement of CRAF1, a relative of TRAF, in CD40 signaling.' cited by the applicant the whole document -----	1-46

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 95/06639

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 30
because they relate to subject matter not required to be searched by this Authority, namely:
"Remark: Although claim 30 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition."
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.